

Character and object*

Jeremy Avigad and Rebecca Morris

September 18, 2012

Abstract

In 1837, Dirichlet proved that there are infinitely many primes in any arithmetic progression in which the terms do not all share a common factor. We survey implicit and explicit uses of *Dirichlet characters* in presentations of Dirichlet's proof in the nineteenth and early twentieth centuries, with an eye towards understanding some of the pragmatic pressures that shaped the evolution of modern mathematical method. We also discuss similar pressures evident in Frege's treatment of functions, and the nature of mathematical objects.

Contents

1	Introduction	2
2	From methodology to metaphysics	3
3	Functions in the nineteenth century	11
3.1	The generalization of the function concept	11
3.2	Other aspects of the evolution of the function concept	17
4	Dirichlet's theorem	19
4.1	Euler's proof that there are infinitely many primes	19
4.2	Group characters	21
4.3	Dirichlet characters and L -series	23
4.4	Contemporary proofs of Dirichlet's theorem	24
5	Modern aspects of contemporary proofs	26
5.1	The hermeneutics of proof	26
5.2	Characters as objects	27

*This essay draws extensively on the second author's Carnegie Mellon MS thesis [81], which contains, in particular, more detailed analyses of some of the proofs of Dirichlet's theorem. We are especially grateful to Michael Detlefsen and the participants in his *Ideals of Proof* workshop, which provided helpful feedback on portions of this material in July, 2011. Avigad's work has been partially supported by National Science Foundation grant DMS-1068829 and Air Force Office of Scientific Research grant FA9550-12-1-0370.

6	Dirichlet’s original proof	30
7	Later presentations	35
7.1	Dedekind	35
7.2	de la Vallée-Poussin	41
7.3	Hadamard	44
7.4	Kronecker	46
7.5	Landau	51
8	Mathematical concerns	54
9	Function and object in Frege	63
10	Foundational concerns	68
11	Conclusions	75

1 Introduction

Historians commonly take the “modern” age of mathematics to have begun in the nineteenth century. But although there is consensus that the events of that century had a transformational effect on mathematical thought, it is not easy to sum up exactly what changed, and why. Aspects of the transformation include an increasingly abstract view of mathematical objects; the rise of algebraic methods; the unification of disparate branches of the subject; evolving standards of rigor in argumentation; a newfound boldness in dealing with the infinite; emphasis on “conceptual” understanding, and a concomitant deemphasis of calculation; the use of (informal) set-theoretic language and methods; and concerns to identify a foundational basis to support the new developments.¹ It is still an important historical and philosophical task to better understand these components, and the complex interactions between them.

A good deal of attention has been given to early appearances of set-theoretic and structural language, including the use of equivalence relations and ideals in algebra; the expansion of the function concept in analysis, and its generalization to other mathematical domains; foundational constructions of number systems from the natural numbers to the reals and beyond; and the overall conception of mathematics as the study structures and spaces, often characterized in set-theoretic terms [41]. These provide relatively clear and focused manifestations of the changes that took place.

In this essay, we will consider certain functions, known as “Dirichlet characters,” and their role in proving a seminal 1837 theorem due to Dirichlet, which states that there are infinitely many prime numbers in any arithmetic progression in which the terms do not all share a common factor. As far as abstract objects go, characters are fairly benign. For a given positive integer m , there are

¹The essays in Ferreirós and Gray [42] provide an overview.

only finitely many characters modulo m , and each one can be described exactly by giving its value on the finitely many residue classes modulo m . Moreover, it is not terribly hard to provide a concrete and exhaustive description of the set of all such characters. Nonetheless, we will argue that the evolution of the treatment of characters over the course of the nineteenth century illustrates important themes in the overall transformation of mathematical thought. Indeed, we will try to call attention to changes in mathematical method that have by now become so ingrained that it is hard for us today to appreciate their significance.

This paper can be read in different ways. It can be viewed, narrowly, as a study of the history of Dirichlet's theorem, with an attempt to understand the pragmatic pressures that shaped that history. More grandiosely, it can be viewed as an attempt to come to terms with the nature of mathematical method and the objects of mathematical study. Along the way, we make some tentative attempts to integrate the narrow and grandiose views.

The first such attempt begins in the next section where, to provide philosophical context, we consider traditional metaphysical concerns in the philosophy of mathematics, and the ways our case study bears on these concerns. In Section 3, we provide historical context as well, by situating our study amidst a host of topics related to the development of the modern function concept. In Section 4, we discuss contemporary presentations of Dirichlet's proof, and in Section 5, we highlight various senses in which these presentations treat characters as objects. In contrast, Section 6 describes Dirichlet's own presentation of his proof, in which the notion of a character does not figure at all. Section 7 then traces a gradual transition, as characters are transformed from shade-like entities in the original proof to the fully embodied objects we take them to be today. Section 8 analyzes the forces that shaped the transition. We return, in Section 9, to metaphysical considerations, as we consider Frege's conflicted attitudes towards the treatment of functions as objects. In Section 10, we argue that there are important commonalities between Frege's foundational concerns and mathematical concerns regarding the proper treatment of functions. Finally, in Section 11, we bring the various strands of our narrative to a point of closure.

2 From methodology to metaphysics

The philosophy of mathematics has long been concerned with the nature of mathematical objects, and the proper methods for gathering mathematical knowledge. But as of late some philosophers of mathematics have begun to raise questions of a broader epistemological character: What does it mean to properly *understand* a piece of mathematics? In what sense can a proof be said to *explain* a mathematical fact? In what senses can one proof be viewed as better than another that establishes the same theorem? What makes a concept fruitful, and what makes one definition more natural or appropriate than another? Why are certain historical developments viewed as important advances?

Questions like these are sometimes classified as pertaining to the *methodology* of mathematics, in contrast to more traditional metaphysical concerns. In this section, we will argue that this distinction is unfortunate, in that a substantive metaphysics has to engage with methodological considerations. In particular, we will explain the sense in which we take it that the methodological study we undertake below bears on a metaphysical understanding.

Let us start by distinguishing between two kinds of questions one can ask, having to do with the existence of mathematical objects. On the one hand, we can ask question such as:

- Is there a nontrivial zero of the Riemann zeta function whose real part is not equal to $1/2$?
- Are there noncyclic simple groups of odd order?

These are fundamentally *mathematical* questions. Answering them is not easy: the Riemann hypothesis posits a negative answer to the first, while the Feit-Thompson theorem, a landmark in finite group theory, provides a negative answer to the second. But even in the first case, where we do not know the answer to the question, we feel that we have a clear sense as to what kind of argument would settle the issue one way or another. Put simply, questions like these can be addressed using conventional mathematical methods. In contrast, there are questions like these:

- Do the natural numbers (really) exist, and what sorts of things are they?
- Are there infinite totalities?
- What kind of sets and functions exist (if any), and what properties do they have?
- Are there infinitesimals, fluxions, fluents, and ultimate ratios?

These are questions as to the ultimate nature of mathematics and its objects of study, and seem to call for a more general, open-ended *philosophical* analysis.

The distinction between the two types of questions may call to mind the logical positivists' distinction between questions that are “internal” to a linguistic framework, and “external” or “pragmatic” questions pertaining to the choice of a framework itself. Some take this distinction to have been repudiated, decisively, by the criticisms of W. V. O. Quine [85]. But keep in mind that Quine's arguments, which were directed against the claim that there is a sharp, principled distinction between the two sorts of questions, were not meant to show that there is no difference between them at all. In locating both kinds of questions on the common continuum of scientific inquiry, he did not deny that different kinds of questions call for different sorts of answers; indeed, his influential *Word and Object* [88] is an extended exploration of the considerations that he took to bear on “philosophical” questions of the latter sort. Nothing we say below commits us to a sharp distinction, and it seems relatively uncontroversial to say that insofar as any rational arguments can be brought to bear on the

second group of questions, these will look different from the kinds of arguments that are brought to bear on the first.

So how shall we go about trying to answer them? Let us consider two approaches.

The first is to turn to a foundational theory, like set theory, for answers. Fixing a definition of the natural numbers provides an answer to the first of the more philosophical questions, and, at the same time, a positive answer to the second. Sets and functions are then understood to have (at least) the properties that the axioms of set theory say they have, and Leibniz' infinitesimals and Newtons' fluxions, fluents, and ultimate ratios can be interpreted in various ways, all perfectly well understood. In this way, foundational reduction serves to reduce philosophical questions to properly mathematical ones, by interpreting them in straightforward mathematical ways.

Foundational reduction of this sort has been one of the most significant contributions of philosophical inquiry to the sciences, providing mathematics with a uniform language for constructing new objects, and clear means of investigating their properties. One might object to this approach to answering our "philosophical questions" on the grounds that there are multiple foundational theories on offer. After all, it would be disappointing if adjudicating ontological questions required obtaining consensus as to whether, for example, set-theoretic foundations or category-theoretic foundations should have the final say. Moreover, even a particular choice of foundation leaves us with too many options; once again, it would be disappointing if our ontological deliberations required us to come to a consensus as to whether the natural numbers are "really" von Neumann ordinals or elements of some other isomorphic structure. But foundational frameworks tend to be largely inter-interpretable,² and the structural nature of mathematics tempers the problem of the overspecificity of a particular choice of definitions. We can characterize the natural numbers uniquely up to isomorphism, and know full well that for most purposes it makes no difference which structure we choose to represent them. With caveats like these, foundational reduction does quite a good job at what it was designed to do.

We may nonetheless feel that our philosophical questions are trying to get at something that foundational reduction fails to address. That is, we may view them as asking whether, in some broad sense, we are justified in admitting the entities in question into our scientific discourse, and attributing those properties to them that our foundational theories are supposed to confirm. In that respect, set-theoretic reduction only pushes the problem back to asking whether we are justified asserting the existence of sets with the properties that our axioms say they have, and likewise for category-theoretic reduction and other alternatives. To avoid appealing to innate or supernatural acquaintance with the objects

²To be sure, it can be hard to interpret a nonconstructive theory in a constructive one, and there some who reasonably hold that, to be meaningful, mathematics *should* be constructive. So there are substantial foundational issues that are still subject to debate, and we do not mean to suggest otherwise. Indeed, as will become clear below, we believe that the approach we describe in this essay does have bearing on issues relating to the tension between constructive and nonconstructive methods in mathematics.

posited by our foundational theories, we then find ourselves pushed to reverse course and justify those theories by the fact that they give us a uniform and useful account of mathematical objects that, antecedently, we have good reasons to be committed to. And those reasons seem to be exactly the sort of thing that our philosophical questions aim to uncover.

Another shortcoming of foundational reduction is a lack of specificity. By aiming for flexibility and universality, foundational frameworks allow us to contrive an endless stream of definitions, the vast majority of them totally useless. In other words, a foundational theory tells us what we are *allowed* to do, not what we *ought* to do. One may therefore interpret our philosophical questions as part of the broader question as to what sorts of mathematical objects ought to be central to our mathematical and scientific discourse. Axiomatic foundations *per se* are not designed to provide this kind of guidance.

Since mathematizing the philosophical questions failed to provide the right sort of answer, we can move to the other extreme, and try to settle them on purely philosophical grounds. For example, in recent years antirealist denials of the existence of abstract mathematical objects have become popular, cast as forms of “nominalism” or “fictionalism.”³ What these positions tend to have in common is that they are decidedly *not* concerned as to whether abstract mathematical objects should be employed by mathematicians, scientists, psychologists, or logicians in their daily discourse; questions like that are viewed as belonging to the realm of methodology, rather than metaphysics proper. Rather than presume to tell mathematicians what they ought to be doing, contemporary metaphysicians focus their efforts on the more fundamental task of supplying a philosophically sound *interpretation* of ordinary mathematical language and determining whether, from a broad philosophical perspective, mathematical objects “really” exist.

There are at least two reasons to be dissatisfied with such an understanding of metaphysics. The first is that, once ties to any pragmatic context have been severed, only vague intuitions and aesthetic judgments can be brought to bear on the fundamental questions. These can pull in opposite directions. For example, some take it as a fundamental tenet of naturalism that existence claims in the sciences should be taken at face value; since seven is a prime number, prime numbers, and hence numbers, exist. Other naturalists wield Occam’s razor as their weapon of choice, and prefer, in the name of science, to discount scientists’ casual existence claims. One man’s meat is another man’s poison; John Burgess dismisses the act of taking back one’s scientific claims in one’s more philosophical moments as doublethink, while denouncing Occam’s razor as “medieval superstition” [10, p. 39]. But rhetorical flourishes like these are unlikely to settle the issue as to whether a minimal ontology is worth a discordant science. Indeed, Maddy [76] has argued that realism and antirealism are, inherently, equally tenable, and Balaguer [7] has argued that, moreover, there is no fact of the matter as to which of them is true. This seems to be a serious drawback to the abstractly philosophical approach, namely, that it

³See, for example, Leng [71].

reduces us to philosophical arguments as to whether there is really anything at stake. This is not to deny that intuitions and aesthetic judgments have bearing on the acceptability of a metaphysical theory, but rather, to emphasize that intuitions and aesthetics, on their own, are not enough: there has to be some *content* to the theory as well.

There is a further reason to ask more from metaphysics. We have rejected a wholesale internalization of the subject as failing to provide useful insight as to *which* mathematical objects are worthy of our attention, and why. The purely philosophical approach fares even worse in this respect: faced with the question as to which *particular* mathematical objects exist, the realist tells us simply that they all do, while the anti-realist tells us that none of them do. In the end, neither view has anything to say about which objects matter to us, and why.

The state of affairs leaves one longing for the days when metaphysics mattered. From Aristotle to Descartes, and then to Frege, Russell, and early twentieth century philosophers of mathematics, the principal philosophical task was to determine the proper language and method of mathematics, including questions as to what objects we should admit into our ontology and how we should reason about them. In this tradition, contemporary metaphysics should provide us with better means of evaluating our methodological choices, and weighing the pros and cons of our decisions. Despite their different characterizations of the philosophical project, Carnap and Quine share the view that metaphysical questions come down to pragmatic questions as to the choice of a conceptual framework. Here is what Carnap has to say about our scientific commitments to abstract objects:

The acceptance cannot be judged as being either true or false because it is not an assertion. It can only be judged as being more or less expedient, fruitful, conducive to the aim for which the language is intended. Judgments of this kind supply the motivation for the decision of accepting or rejecting the kind of entities.⁴ [14, p. 250]

Quine offers the following amendment:

Consider the question whether to countenance classes as entities. This, as I have argued elsewhere, is the question whether to quantify with respect to variables which take classes as values. Now Carnap has maintained that this is a question not of matters of fact but of choosing a convenient language form, a convenient conceptual scheme or framework for science. With this I agree, but only on the proviso that the same be conceded regarding scientific hypotheses generally. [85, p. 43]

We take these views seriously here, seeing it as an important metaphysical task to clarify the role of our ontological posits with respect to ordinary mathe-

⁴Here and below, when the bibliographic entry of a work includes a reprinted version, page numbers in the references refer to the reprinted version. Similarly, when the bibliographic entry includes an English translation, our translations are taken from that source, unless we indicate otherwise. Where no translation is listed, the translations are our own.

mathematical activity, and evaluate their efficacy towards achieving our mathematical goals. This amounts to something like the naturalist approaches to the philosophy of mathematics as advocated by Kitcher [60], Burgess [12], and Maddy [75], focused on specific aspects of mathematical practice. The task is by no means easy, not least since it is not at all clear how to characterize our mathematical goals or evaluate our success at achieving them. But philosophy should not shy away from hard work.

On this view, metaphysics becomes a kind of means-ends analysis. It is therefore important to keep in mind that scientific ends are usually best described relative to a scientific context. While a mathematician may have good reason to admit certain abstract objects with all their attendant properties, a logician may have good reason to adopt a more minimal ontology when it comes to studying the metamathematical effects of our axiomatic assumptions; and there may be good reasons for psychologists to avoid reference to abstract objects altogether in an account of how children learn to grasp mathematical concepts. It seems fruitless to grant any of these contexts the final say as to whether mathematical objects “really” exist; the objects in question play distinct roles in distinct theories with distinct explanatory and predictive goals. In situations where the scientific contexts overlap, some translation and reinterpretation is inevitable, and part of the metaphysician’s task is to help keep these cross-disciplinary collaborations running smoothly. Our view does not rule out hope for an overarching frame that is broad enough to fit together the various scientific snapshots; but we do insist that, ultimately, the success of such a metaphysics needs to be judged in terms of its contribution to the pragmatic needs of the working scientist.

In this essay, we will focus on the way that ontological commitments bear on distinctly mathematical concerns, supporting us in or hindering us from achieving our mathematical goals. How, then, should such an analysis proceed?

It is instructive to consider those historical situations in which the mathematical community faced possibilities for methodological or ontological expansion and reacted accordingly. For example, it is helpful to consider the ancient Greek idealizations of number and magnitude, and the theory of proportion; the gradual acceptance of negative numbers, and then complex numbers, in the Western tradition; the use of algebraic methods in geometry, infinitesimals in the calculus, points at infinity in projective geometry; the development of the function concept from Euler to modern times; the gradual set-theoretic treatment of algebraic objects like cosets, ideals, equivalence classes in the nineteenth century; and so on. By studying the historical concerns regarding these expansions as well as the pressures that led to their ultimate acceptance, we can hope to better understand the metaphysical considerations that can be brought to bear.

Indeed, at junctures like these, historical developments tend to follow a common pattern. First, expansions are met with resistance, or at least, extreme caution. Sometimes, the expansions can be explained in terms of the more conservative practice; for example, complex numbers can be interpreted as ordered pairs, algebraic solutions to geometric problems can be reinterpreted geometri-

cally, equations can be rewritten to avoid consideration of negative quantities. In other cases, the expansions are not generally conservative, but, at least, can be explained away in particular instances; for example, arguments involving infinitesimals can sometimes be interpreted in terms of “ultimate ratios” in a geometric diagram, and operations on abstract objects can sometimes be understood as operations on explicit representations. This makes it possible to adopt the expansions, tentatively, as convenient shorthand for more tedious but conservative arguments. Over time, the rules and norms that govern the expansions are clarified, and the expansions themselves prove to be convenient, or even indispensable, while they do not cause serious problems. Over time, the mathematical community grows used to them, to the point where they become part of the usual business of mathematics.

Whiggish narratives tend to dismiss such historical hand-wringing and shilly-shallying as short-sighted conservatism on road to progress. We, however, prefer to view it as a rational response to the proposed expansions, whereby the benefits are carefully weighed against the concerns. In hindsight, we tend to make too little of the pitfalls associated with an ontological or methodological expansion. To start with, there are concerns about the *consistency* and *coherence* of the new methods, that is, worries as to whether the changes will lead to mistakes, false results, or utter nonsense, perhaps when employed in situations that have not even been imagined. Kenneth Manders has also emphasized the importance of maintaining *control* of our mathematical practices [78]. Mathematics requires us to be able to come to agreement as to whether a proof is correct, or whether a given inference is valid or not. If new objects come with rules of use that are not fully specified, or vague, or unclear, the practice is in danger of breaking down. In a sense, this concern comes prior to concerns of consistency: if it is not clear what properties abstract magnitudes, negative numbers, complex numbers, infinitesimals, sets, and “arbitrary” functions have, it doesn’t even make sense to ask whether using them correctly will lead to contradictions.⁵

And then there are further concerns as to whether the new methods are *meaningful* and *appropriate* to mathematics. Even if a body of methods is consistent and clearly specified, it may still fail to provide us with the results we are after. If you expect an existence proof to yield certain kinds of information about the object that is asserted to exist, methods that fail to provide that sort of information do not constitute mathematics—or, at least, not the kind of mathematics we should be doing. If you expect a mathematical theory to make scientific predictions that we can act on rationally, it is a serious concern as to whether the new methods can deliver.

In short, the concerns are not easily set aside. What, then, are the factors that might sway a decision in favor of an expansion? Mathematicians tend to wax poetic in their praise of conceptual advances, highlighting the power of new methods, the elegance and naturality of the resulting theory, and the insight and

⁵As Wilson [101] and [97] emphasize, however, mathematics often gets by surprisingly well with concepts that are problematic, incompletely specified, and not fully understood.

depth of the associated ideas. Part of our goal here is to de-romanticize these virtues and gain clarity as to what might be achieved. In many instances, the virtues in question seem to have a lot to do with efficiency and economy of thought:⁶ we tend to value methods that make it possible to solve problems that were previously unsolvable, or simplify proofs and calculations that were previously tedious, complex, and error-prone. Below we will consider specific ways in which ontological and methodological expansions help us manage complex tasks by suppressing irrelevant detail, making key features of a problem salient, and keeping key information ready-to-hand. We will also try to understand the way they make it possible to generalize and extend results, and facilitate the transfer of ideas to other domains.

To summarize our high-level historical model: when mathematics is faced with methodological expansion, benefits such as simplicity, generality, and efficiency are invariably weighed against concerns as to the consistency, cogency, and appropriateness of the new methods. Sufficient benefit encourages us to entertain the changes cautiously, while trying to minimize the dangers involved. Cogency is obtained by working out the norms and conventions that govern the new methods. Consistency may not be guaranteed, but our experiences over time can bolster our faith that the new methods do not cause problems. In this regard, initial checks that the new methods are partially conservative over the old ones helps preserve mathematical meaning, and reassures us that even if the new methods turn out to be problematic, one will be able to restrict their scope in such a way that preserves their utility.⁷ A metaphysics of mathematics should give us better means to evaluate such expansions: to talk about cogency of a mathematical argument and whether it delivers the desired content; and to understand the ways in which our ontological posits and methodological expansions improve our ability to reason effectively.

Philip Kitcher invoked a similar model in his important book, *The Nature of Mathematical Knowledge* [59]. Chapter 9, “Patterns of mathematical change,” discusses the historical tendency to weigh “problem-solving dividends” against the perceived costs of an extension, which stand “on the other side of the ledger” [59, p. 199]. Kitcher’s case studies, which include the introduction of Cartesian methods to geometry and nineteenth century developments in analysis, highlight considerations very similar to the ones we discuss here. One small difference is that Kitcher focused on foundational developments in which the methods under consideration make it possible to solve problems that were previously unsolvable while, at the same time, posing clear problems, such as false conclusions and uncomfortable gaps in reasoning. Below, we focus on evolving presentations of a proof whose correctness was never in question, showing that such cost-benefit considerations bear upon everyday mathematical developments as well. A more striking difference is that Kitcher invokes his historical model in support of an empiricist epistemology of mathematics that runs counter to what he calls the

⁶The phrase is borrowed from Ernst Mach’s *The Science of Mechanics* [73]; we are grateful to Michael Detlefsen for bringing this to our attention

⁷Wittgenstein’s discussion of contradiction is interesting in this regard; see [105, Lectures XI–XII].

traditional “apriorist program.” His concerns trace back to the Carnap-Quine debate as to whether one can, and should, maintain some sort of principled distinction between analytic and synthetic knowledge. These issues are largely orthogonal to the concerns addressed here: although our case study can support a better understanding of the kinds of rational considerations that bear upon mathematical developments, it is not wedded to any particular stand on the analytic / synthetic distinction.

With this long-winded discussion behind us, we can now state our goals more clearly. In this essay, we will explore the evolution of the mathematical treatment of certain types of functions in nineteenth century mathematics, from an appropriately metaphysical point of view.

3 Functions in the nineteenth century

3.1 The generalization of the function concept

Nineteenth century mathematicians dealt with a number of objects that we, today, view as instances of the function concept. Analysis dealt with functions defined on the continuum, that is, functions from the real numbers, $\mathbb{R}$, to $\mathbb{R}$. Cauchy and others extended the subject to include functions from the complex numbers, $\mathbb{C}$, to $\mathbb{C}$. Analysts also commonly dealt with sequences and series, which can be viewed as functions from the natural numbers, $\mathbb{N}$, to a target domain, such as the integers or the reals. Number theory dealt with functions like the Euler totient function, φ , which maps $\mathbb{N}$ to $\mathbb{N}$. Geometry dealt with transformations of the Euclidean or projective plane, which can be viewed as invertible maps from the space to itself. Galois’ theory of equations focused on substitutions, or permutations, of a finite set A , which is to say, bijective functions from A to A . Below we will see how developments in number theory led to the study of characters, that is, functions from the integers, $\mathbb{Z}$, to $\mathbb{C}$, or from the group $(\mathbb{Z}/m\mathbb{Z})^*$ of units modulo m to $\mathbb{C}$. By the end of the century, Dedekind and Cantor considered arbitrary mappings, or correspondences, between domains.

Despite this diversity, most of the literature on the evolution of the function concept (including [16, 62, 72, 79, 107]) has focused on functions on the real or complex numbers. Surveys typically trace the evolution of the concept from the introduction of the term “function” (*functio*) by Euler [36] in 1748, to the mention of “arbitrary functions” (*fonction arbitraire*) in the title of Dirichlet’s seminal paper of 1829 [25], to the dramatically opposed treatment of the function concept by Riemann and Weierstrass.⁸ There is a very good reason for this: in the early nineteenth century, the word “function” was used almost exclusively in this narrow sense. Indeed, there is little in the early nineteenth century literature that suggests any family resemblances between the kinds of

⁸See Bottazzini and Gray [52] for a history of complex function theory, which includes a detailed treatment of the work of Riemann and Weierstrass, in particular.

mathematical objects enumerated in the last paragraph, let alone subsumption under a single overarching concept.

For example, sequences of real numbers were referred to as sequences or series, or often just introduced by displaying the initial elements $a_0, a_1, a_2, \dots$ with an ellipsis at the end. The words “series” (*série* in French, *Reihe* in German) was also used to describe a finite or infinite sum. For example, Dirichlet wrote the following in 1835:

Now let

$$F(\alpha) = b_0 + b_1 \cos \alpha + b_2 \cos 2\alpha + \dots = \sum b_i \cos i\alpha$$

be an arbitrary finite or infinite series, whose coefficients are independent of α .⁹ [26, p. 249]

The dependence of the coefficient b_i on i was indicated by the subscript, just as we do today.

Terminology governing number-theoretic functions was more varied. In the eighteenth century, Euler tended to refer to number-theoretic functions as “symbols” or “characters.” For example, when introducing what we now call the totient function in 1781, he wrote:

... let the character πD denote that multitude of numbers which are less than D and which have no common divisor with it.¹⁰ [37, p. 2]

In 1801, in §38 of the *Disquisitiones Arithmeticae* [50], Gauss introduced the totient function as a notation:

For brevity we will designate the number of positive numbers which are relatively prime to the given number and smaller than it by the prefix φ . We seek therefore φA .¹¹

In the following section, he referred to the “character” φ . Later, in §52, he defined another number-theoretic function, ψ , and in §53 compared it to the φ “symbol” (*signo*). In the *Disquisitiones*, the word “function” is never used to describe such entities.

Early in the nineteenth century, in projective geometry, Möbius studied general “relations” between figures in the plane, such as “affinities” and “collineations.”¹² In his *Erlangen Program* of 1872, Klein considered invertible “transformations” (*Transformationen*) of a space, and the group they form under the operation

⁹ “Es sei jetzt:

$$F(\alpha) = b_0 + b_1 \cos \alpha + b_2 \cos 2\alpha + \dots = \sum b_i \cos i\alpha$$

eine beliebige endliche oder unendliche Reihe, deren Coefficienten von α unabhängig sind.”

¹⁰ “Quod quo facilius praestari possit, denotet character φD multitudinem istam numerorum ipso D minorum, et qui cum eo nullum habeant divisorem communem.” [37, p. 19]

¹¹ “Designemus brevitatis gratia multitudinem numerorum positivorum ad numerum datum primorum ipsoque minorum per praeifixum characterem φ . Quaeritur itaque φA .”

¹² See the discussion in Wussing [106, pp. 35–40].

of “composition” (*Zusammensetzung*) [61]. Once again, in neither case is there any hint that such transformations bear a connection to a more general function concept.

In algebra, Galois’ groups consisted of “substitutions” (*substitutions*) of the roots. Cayley’s famous 1854 paper, which provided the first axiomatization of the group concept, begins as follows:

Let θ be a symbol or operation, which may, if we please, have for its operand, not a single quantity x , but a system $(x, y, \dots)$, so that

$$\theta(x, y, \dots) = (x', y', \dots),$$

where $x', y', \dots$ are any functions whatever of $x, y, \dots$, it is not even necessary that $x', y', \dots$ should be the same in number with $x, y, \dots$. In particular, $x', y', \dots$ &c. may represent a permutation of $x, y, \dots$, θ is in this case what is termed a substitution; and if, instead of a set $x, y, \dots$, the operand is a single quantity x , so that $\theta x = x' = fx$, θ is an ordinary function symbol.¹³ [15, p. 123]

Notice that although we can assume that each quantity $x', y', \dots$ is a quantity of $x, y, \dots$, for Cayley, θ is, in general, an operation and not a function. As examples of cases where θ is a function, he singled out multiplication by quaternions and examples arising from the study of elliptic functions. But he makes it clear that it is the operations, more generally, that are assumed to satisfy the associative law, under composition:

A symbol $\theta\varphi$ denotes the compound operation, the performance of which is equivalent to the performance, first of the operation φ , and then of the operation θ ; $\theta\varphi$ is of course in general different from $\varphi\theta$. But the symbols $\theta, \varphi, \dots$ are in general such that $\theta.\varphi\chi = \theta\varphi.\chi$, &c., so that $\theta\varphi\chi, \theta\varphi\chi\omega$, &c. have a definite signification independent of the particular mode of compounding the symbols...

Dedekind’s later treatment of Galois theory [23] focused on the group of automorphisms of a field rather than permutations of roots, but even in this context he used the term “substitution” for an invertible map from a field to itself, and “permutation” for a substitution that is moreover a homomorphism with respect to the field structure. Once again, there seems to be nothing to link these substitutions and permutations with the function concept from analysis. Even as late as 1895, when Cantor presented his mature theory of the infinite [13], he described a one-to-one correspondence as a “law of association” (*Zuordnungsgesetz*) between sets, with nothing to suggest that these correspondences had anything to do with the objects of function theory with which he had begun his mathematical career.

The shift from using the term “function” exclusively for functions defined on the continuum to those defined on more general domains was gradual. We

¹³This excerpt is quoted and discussed in Pengelley [84].

have found a very early instance of the phrase “number-theoretic function” (*zahlentheoretische Funktion*) in an 1850 paper by Eisenstein, which begins with a notably self-conscious justification for the use of the term:

Since, with the concept of a function, one moved away from the necessity of having an analytic construction, and began to take its essence to be a tabular collection of values associated to the values of one or several variables, it became possible to take the concept to include functions which, due to conditions of an arithmetic nature, have a determinate sense only when the variables occurring in them have integral values, or only for certain value-combinations arising from the natural number series. For intermediate values, such functions remain indeterminate and arbitrary, or without any meaning.¹⁴ [35, p. 706]

We might have expected Eisenstein to observe, more simply, that once one starts to think of a function as a correspondence between input and output values, it is reasonable to transfer the notion to correspondences between domains other than the real or complex numbers, described by means other than an analytic expression. Instead, he takes the surprising tack of viewing number-theoretic functions as *partial functions on the real numbers*. This is a nice illustration of the fact that concepts are often stretched and transformed in surprising ways, a theme that is the central study of Mark Wilson’s book, *Wandering Significance* [102]. In any case, the passage makes it clear that the modern notion of a function defined on arbitrary domains was far from the mid-nineteenth century mindset.

We have already noted that in 1854, Cayley considered multiplication by a quaternion as a function. This seems to fit with the view of the natural progression from real numbers to complex numbers to quaternions, as generalizations of the concept of magnitude. One finds another hint of expansion in the title of Dedekind’s 1854 *Habilitation* lecture, “On the introduction of new functions in mathematics” (“Über die Einführung neuer Funktionen in der Mathematik”) [20]. In this lecture, Dedekind discussed the way that the domain of natural numbers was gradually expanded to include the integers, real numbers, and complex numbers, while extending and preserving the properties of familiar operations like addition, and division. But the evocative use of the word “function” in the title is tempered by the contents of the lecture itself, where the word “operation” is used exclusively when it comes for functions defined on the natural numbers and integers.

¹⁴ “Seit man bei dem Begriffe der Funktion von der Nothwendigkeit der analytischen Zusammensetzung abgehend, das Wesen derselben in die tabellarische Zusammenstellung einer Reihe von zugehörigen Werthen mit den Werthen des order der (mehrerer) Variablen zu setzen anfang, war es möglich, auch solche Funktionen unter diesen Begriff mit aufzunehmen, welche aus Bedingungen arithmetischer Natur entspringend nur für ganze Werthe oder nur für gewisse aus der natürlichen Zahlenreihe hervorgehende Werthe und Werth-Combinationen der in ihnen vorkommenden Variablen einen bestimmten Sinn erhalten, während sie für die Zwischenwerthe entweder unbestimmt und willkürlich oder ohne alle Bedeutung bleiben.” We are grateful to Wilfried Sieg for help with the translation.

To our knowledge, it is not until 1879 that one finds the word “function” used with any approximation to the modern sense; and then, in that year, it occurred in two remarkable sources. The first is the work of Dedekind: in 1879, in a supplement dealing with quadratic forms in the third edition of his presentation of Dirichlet’s lectures on number theory [29], he defined the general notion of a character on a class group (that is, a function which maps equivalence classes of ideals in an algebraic number field to the complex numbers):

...the function $\chi(\mathfrak{a})$ also possesses the property that it takes the same value on all ideals $\mathfrak{a}$ belonging to the same class A ; this value is therefore appropriately denoted by $\chi(A)$ and is clearly always an h th root of unity. Such functions χ , which in an extended sense can be termed *characters*, always exist; and indeed it follows easily from the theorems mentioned at the conclusion of §149 that the class number h is also the number of all distinct characters $\chi_1, \chi_2, \dots, \chi_h$ and that every class A is completely characterized, i.e. is distinguished from all other classes, by the h values $\chi_1(A), \chi_2(A), \dots, \chi_h(A)$.¹⁵

By 1882, Weber [100] described characters on arbitrary groups as “functions.”¹⁶ In his foundational essay *Was sind und was sollen die Zahlen* [22] of 1888, Dedekind considered arbitrary mappings (*Abbildung*) between domains, but, curiously, seems to distinguish mappings from functions.¹⁷

The other landmark source of 1879 is Frege’s *Begriffsschrift* [43]. Here Frege defined the notion of a function as follows:

If, in an expression (whose content need not be a judgeable content), a simple or complex symbol occurs in one or more places, and we think of it as replaceable at all or some of its occurrences by another symbol (but everywhere by the same symbol), then we call the part of the expression that on this occasion appears invariant the function, and the replaceable part its argument.¹⁸ [43, §9]

¹⁵ “...die Function $\chi(\mathfrak{a})$ ausser der Eigenschaft ... noch die andere besitzt, für alle derselben Classe A angehörenden Ideale $\mathfrak{a}$ denselben Werth anzunehmen, welcher mithin zweckmässig durch $\chi(A)$ bezeichnet wird und offenbar immer eine h^{te} Wurzel der Einheit ist. Solche Functionen χ , die man im erweiterten Sinn *Charaktere* nennen kann, existiren immer, und zwar geht aus den am Schlusse des §149 erwähnten Sätzen leicht hervor, dass die Classenanzahl h zugleich die Anzahl aller verschiedenen Charaktere $\chi_1, \chi_2, \dots, \chi_h$ ist, und dass jede Classe A durch die ihr entsprechenden h Werthe $\chi_1(A), \chi_2(A), \dots, \chi_h(A)$ vollständig charakterisirt, d.h. von allen anderen Classen unterschieden wird.” The quotation appears in §178 in the 1879 edition of the *Vorlesungen* [29], and in §184 of the 1894 edition, which is reproduced in Dedekind’s *Werke* [24]. The translation above is by Hawkins [56, p. 149].

¹⁶We discuss Weber’s 1882 paper, and provide more background on the history of characters, in Section 7.1, below.

¹⁷See §135 in [22]. The relationship between the two notions will be discussed in forthcoming work by Wilfried Sieg and the second author.

¹⁸“Wenn in einem Ausdrucke, dessen Inhalt nicht beurtheilbar zu sein braucht, ein einfaches oder zusammengesetztes Zeichen an einer oder an mehrern Stellen vorkommt, und wir denken es an allen oder einigen dieser Stellen durch Anderes, überall aber durch Dasselbe ersetzbar, so nennen wir den hierbei unveränderlich erscheinenden Theil des Ausdruckes Function, den ersetzbaren ihr Argument.”

While this definition is couched in terms of expressions, Frege later formulated an account that does not depend on linguistic apparatus. We will consider this account in more detail in Sections 9 and 10, but make a few brief remarks here. In 1891, Frege noted that his conception of function extended previous ones in two ways: by enlarging the collections of signs that could be used to construct a functional expression, and by enlarging the domain of possible arguments for functions [45].¹⁹ Regarding the first extension, Frege allowed signs such as the equality symbol to occur in functional expressions, thus allowing “ $x^3 + x^2 + x + 1 = 0$ ” to be classified as a functional expression. Regarding the second extension, Frege wrote:

Not merely numbers, but objects in general, are now admissible; and here persons must assuredly be counted as objects.²⁰ [45, p. 17]

The syntax of Frege’s logical language allows, moreover, for higher-order functionals, which is to say, functions which take elements of a domain of functions as arguments. Although Frege was influenced by Riemann and developments in the theory of complex functions [95], we will see in Section 9 that some features of his treatment of the function concept distinguish it from the contemporary mathematical notion. Nonetheless, in the context of the history we have sketched here, Frege’s dramatic expansion was novel and bold.

Today, we are apt to look back at the history and wonder what mathematicians before Dedekind and Frege were missing, that is, why they didn’t realize the extent to which they could simplify terminology and notation and eliminate conceptual clutter. After all, all they had to do was write $f : A \rightarrow B$ to denote a function f between two arbitrary domains, A and B , and recognize sequences, number-theoretic functions, permutations, and transformations as instances of such. But this seems to us to be the wrong question to ask. A better one is this: why would anyone want to view these patently different entities as instances of a single concept? In practice, they are described in very different ways: analytic functions were typically given by piecewise analytic expressions; number theoretic functions were given by implicit algorithms, such as counting the numbers less than and relatively prime to some number; Galois’ substitutions were given by explicit lists; early geometric transformations were given by geometric constructions. Moreover, the different kinds of objects support very different operations: functions in analysis can sometimes be composed, but sequences and series cannot; substitutions (and Cayley’s “operations”) can always be inverted, whereas number-theoretic functions generally cannot; one can sum an expression over all the permutations of a finite set, while one certainly cannot sum an expression over all the functions from $\mathbb{R}$ to $\mathbb{R}$; notions of continuity and differentiability make no sense outside the realm of analysis; and so on.

As noted in Section 2, there are startup costs involved in fashioning coherent means of dealing with new abstract objects, and significant concerns. Thus one

¹⁹See [8, pp. 137 and 140].

²⁰“Es sind nicht mehr bloß Zahlen zuzulassen, sondern Gegenstände überhaupt, wobei ich allerdings auch Personen zu den Gegenständen rechnen muß.”

should not expect to see an investment in the unification of the function concept until there was either a pressing need or clear benefits to be had. Our study below will suggest that some of these benefits stem from the uniform treatment of algebraic structures composed of functions. For example, the set of automorphisms of a structure form a group under composition; but there is little reason to recognize permutations, geometric transformations, or automorphisms of a field as such until one has particularly useful things to say about groups of automorphisms in general. Similarly, one can view a number of common mathematical constructions as quotienting by the kernel of a suitable function, but there is no point to making the effort to recognize them as such until one realizes that substantial things can be said about the commonality. Alternatively, it is also possible that the unification of the function concept is best viewed as a by-product of the drive towards subsuming mathematical objects under a single unifying foundation, a move which, in turn, might have been encouraged by other mathematical needs.

It is beyond the scope of our study to speculate more than this as to the factors that encouraged the development of a general concept of function and led to its gradual acceptance. But noting the absence of an overarching function concept in the early nineteenth century serves to clarify the nature of the project here. When we consider the history of number-theoretic characters in proofs of Dirichlet's theorem, we are studying the evolving mathematical treatment of *what we now take to be* certain kinds of functions, in the hopes that understanding the forces that guided that evolution will help illuminate the forces that shaped the evolution of modern mathematics more generally. In doing so, we will focus on the way characters were used and the properties that were ascribed to them, while, for the most part, steering clear of the question as to how and why they came to be viewed as instances of the general function concept.

3.2 Other aspects of the evolution of the function concept

We have seen that it was not until the latter half of the nineteenth century that there was any hint of a general notion of a function between any two domains, and that it was a while before what we currently take to be instances of the function concept were subsumed under such a general notion. Let us label this trend the *generalization* (or, perhaps, *unification*) of the function concept. The point of this section is to note that this is not the only sense in which the function concept evolved over the course of the century.

To start with, there was also the *extensionalization* of the function concept. Recall that, in Euler's time, a function from the real numbers to the real numbers was expected to have a representation in terms of a certain kind of analytic expression. Arguments involving functions could then presuppose such a representation. Precisely this feature of Cauchy's treatment of Fourier series was the target of Dirichlet's criticism in the opening paragraphs of his 1829 paper [25] on that same subject. Cauchy, like Dirichlet, was concerned with the question as to when the Fourier series of a real-valued function converges to the value of

the original function at a given point. But Cauchy’s analysis presupposed that the function in question is represented by a power series, which could then be applied to complex arguments. The use of the phrase “arbitrary function” in Dirichlet’s title signaled his intention to lift this restriction. Even though the paper dealt primarily with continuous functions, the hypothesis of continuity was expressed purely extensionally, which is to say, in terms of the values that a function takes at its arguments. In other words, Dirichlet took great pains not to treat analytic operations like differentiation as symbolic operations, but, rather, as operations on the functions themselves, viewed extensionally. As Dirichlet famously emphasized, this renders the concept of function open-ended, in that all that is need is some determinate relationship between input and output values. In particular, this made it possible for him to consider, as a difficult example, functions which take one value on the rationals, and another value on the irrational numbers.

This brings us to another aspect of the evolution of the notion of a function, which we will call the *liberalization* of the function concept. Setting aside issues of unification and generality, there is also the issue of the language and methods one helps oneself to in *defining* functions in a particular domain. As Frege later noted,²¹ the very act of accepting a definition by cases on whether an argument is rational or not was a striking move on Dirichlet’s part. This paves the way to the use of more dramatically non-constructive methods in describing functions from the reals to the reals, say, in terms of limits and other infinitary operations. Riemann’s contributions to function theory were especially novel in this regard. Some, like Dedekind, hailed the fact that Riemann’s methods made it possible to consider functions without having a particular representation to work with, or method of computation. Others, like Weierstrass, held the Riemannian approach to be defective for just this reason.²²

Finally, there is the issue of the *reification* of the function concept, which is to say, the decision to treat functions as objects, on par with mathematical objects like the natural numbers. In Section 5, we will have a lot more to say about what this amounts to.

We have already indicated that we will have little more to say about the generalization of the function concept. And, since characters are fairly simple combinatorial objects, we can avoid issues having to do with liberal uses of the infinite. We will be quite concerned, however, with issues regarding the reification of characters, and their treatment as extensional objects. Thus our case study considers some important aspects of the evolution of the function concept, but ignores others, as well as the broader question as to how all the various components are related to one another.

²¹See the discussion in Section 8.

²²See Bottazzini and Gray [52]. For an interesting exploration of the ways that nineteenth century analysis expanded to incorporate a more liberal understanding of the function concept, see Chorlay [16].

4 Dirichlet's theorem

Two integers, m and k , are said to be *relatively prime*, or *coprime*, if they have no common factor. In 1837, Dirichlet proved the following:

Theorem 4.1. *If m and k are relatively prime, the arithmetic progression $m, m + k, m + 2k, \dots$ contains infinitely many primes.*

In other words, if m and k are relatively prime, there are infinitely many primes congruent to m modulo k . Dirichlet pointed out that Legendre assumed this fact, without proof, in 1788 [70], when proving the law of quadratic reciprocity. In his *Disquisitiones Arithmeticae*, Gauss noted this gap in Legendre's work after presenting his own proof of the law of quadratic reciprocity, one which does not rely on Theorem 4.1.²³ But nor did Gauss ever provide a proof of that theorem. Dirichlet's own proof is striking, not only due to the fact that it finally established Legendre's conjecture, but also due to its sophisticated use of the methods of analysis in establishing a purely number-theoretic assertion. Dirichlet noted [27, pp. 309–310] that his method was inspired by a proof due to Euler [36, Chapter XV] that there are infinitely many primes, though Dirichlet's ideas go considerably beyond Euler's own. In this section, we will describe Euler's proof, and then define the modern notion of a group-theoretic character, which supports the generalization to arbitrary characters. We will then describe contemporary proofs of Dirichlet's theorem, using modern terminology and notation, as presented in textbooks such as Everest and Ward [38, pp. 207–224]. In Section 5, we will use this presentation to frame our discussion of the historical development, which occurs in Sections 6 and 7.

There is a sense, however, in which our modern presentation is old-fashioned. The use of m and k for the initial value and common difference in the statement of Theorem 4.1 is due to Dirichlet, and was picked up by a number of successive authors, including Dedekind and Hadamard. Modern presentations are more apt to use a and d , but although the use of m and k may feel alien to readers familiar with contemporary textbook proofs, it will facilitate the historical comparisons later on. (Our reason for using the variable q to range over the prime numbers will similarly become clear when we discuss Dirichlet's original proof.)

4.1 Euler's proof that there are infinitely many primes

In the *Elements*, Euclid proved that there are infinitely many primes, but his proof does not provide much information about how they are distributed. Euler, in his *Introductio in analysin infinitorum* [36], proved the following:

Theorem 4.2. *The series $\sum_q \frac{1}{q}$ diverges, where the sum is over all primes q .*

This implies that there are infinitely many primes, but also says something more about their density. For example, since we know that the series $\sum_n \frac{1}{n^2}$ is

²³See Gauss' remarks in [50, §150].

convergent, it tells us that, in a sense, there are “more” primes than there are squares.

Euler’s proof of Theorem 4.2 centers around his famous zeta function,

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s},$$

defined for a real variable s . (The zeta function was later extended by Riemann to the entire complex plane via analytic continuation.) It is not hard to show that the series $\zeta(s)$ converges uniformly on the interval $[a, \infty)$, where a is any number strictly greater than 1. For $s > 1$, the infinite sum can also be expressed as an infinite product:

$$\sum_{n=1}^{\infty} n^{-s} = \prod_q \left(1 - \frac{1}{q^s}\right)^{-1},$$

where the product is over all primes q . This is known as the *Euler product formula*. Roughly, this holds because we can write each term of the product as the sum of a geometric series,

$$\left(1 - \frac{1}{q^s}\right)^{-1} = 1 + q^{-s} + q^{-2s} + \dots$$

and then expand the product into a sum. The unique factorization theorem tells us that every integer $n > 1$ can be written as a product $q_1^{i_1} \cdot q_2^{i_2} \cdots q_k^{i_k}$, and so the term $n^{-s} = q_1^{-i_1 s} \cdot q_2^{-i_2 s} \cdots q_k^{-i_k s}$ will occur exactly once in the expansion. Since we are dealing with infinite sums and products, the Euler product formula implicitly makes a statement about limits, and some care is necessary to make the argument precise; but this is not hard to do.

If we take the logarithm of each side of the product formula and appeal to properties of the logarithm function, we obtain

$$\log \sum_{n=1}^{\infty} n^{-s} = \sum_q -\log \left(1 - \frac{1}{q^s}\right).$$

Using the Taylor series expansion

$$\log(1 - x) = -x - x^2/2 - x^3/3 - \dots$$

and changing the order of summations yields

$$\log \sum_{n=1}^{\infty} n^{-s} = \sum_q \frac{1}{q^s} + \sum_{n=2}^{\infty} \frac{1}{n} \sum_q \frac{1}{q^{ns}}.$$

At this stage, keep in mind that we want to show that $\sum_q \frac{1}{q}$ diverges, and notice that the first term on the right hand side of the above equation is $\sum_q \frac{1}{q^s}$. Thus

we should consider what happens as s tends to 1 from above. One can show that the second term on the right-hand side is bounded by a constant that is independent of s , a fact that can be expressed using “big O” notation as follows:

$$\log \sum_{n=1}^{\infty} n^{-s} = \sum_q \frac{1}{q^s} + O(1). \quad (1)$$

As s approaches 1 from above, the left-hand side clearly tends to infinity. Thus, the right-hand side, $\sum_q \frac{1}{q^s}$, must also tend to infinity, which implies that $\sum_q \frac{1}{q}$ diverges.

4.2 Group characters

Just as Euler’s proof shows that there are infinitely many primes by establishing that the series $\sum_q \frac{1}{q}$ is divergent, proofs of Dirichlet’s theorem establish that there are infinitely many primes q congruent to m modulo k by showing that the series $\sum_{q \equiv m \pmod{k}} \frac{1}{q}$ is divergent. Remember that we are assuming that m and k are relatively prime. The residues modulo k that are, moreover, relatively prime to k form a multiplicative group. To adapt the argument above, we need series that are more refined than the zeta function, and a device that enables us to focus attention on the integers that are congruent to m modulo k . This is where the notion of a group-theoretic character comes in.

Let G be a finite abelian group, written multiplicatively, with an identity denoted by 1. A *character* χ on G is a homomorphism from G to the nonzero complex numbers, $\mathbb{C}^*$. In other words, a character χ is a nonzero function satisfying $\chi(g_1 g_2) = \chi(g_1) \chi(g_2)$ for every g_1 and g_2 in G . Since $\chi(1) = \chi(1 \cdot 1) = \chi(1) \chi(1)$ and $\chi(1)$ is nonzero, we have $\chi(1) = 1$. Since G is a finite group, for every g there is a least $n > 1$ satisfying $g^n = 1$; this n is called the *order* of g and denoted $o(g)$. The fact that $\chi(g)^{o(g)} = \chi(g^{o(g)}) = \chi(1) = 1$ means that $\chi(g)$ is a “root of unity” for every g in G . The character which is equal to 1 for every g in G is called the *trivial character* and denoted by χ_0 .

Define the product $\chi \cdot \psi$ of two characters pointwise, by

$$(\chi \cdot \psi)(g) = \chi(g) \psi(g),$$

for every g in G . This multiplication is commutative, and we have $\chi \cdot \chi_0 = \chi$ for every character χ , which is to say, χ_0 is a multiplicative identity. Recall that if ω is any complex root of unity, its complex conjugate, $\bar{\omega}$, is also a root of unity, satisfying $\omega \bar{\omega} = 1$. We can also lift the operation of conjugation to characters, defining $\bar{\chi}$ by the equation $\bar{\chi}(g) = \overline{\chi(g)}$ for each g . Then clearly we have $\chi \cdot \bar{\chi} = \chi_0$. In other words, the set of characters on G forms an abelian group, with $\chi^{-1} = \bar{\chi}$. We will denote this group $\hat{G}$.

The following theorem is fundamental:

Theorem 4.3. *If G is any finite abelian group, $\hat{G}$ is isomorphic to G . In particular, $|\hat{G}| = |G|$.*

To see this, first consider the case where $G = \langle g \rangle = \{1, g, g^2, \dots, g^{n-1}\}$ is a cyclic group generated by an element g of order n . Then any character χ on G has to map g to an n th root of unity, ω . This determines the behavior of χ completely, since then we have $\chi(g^i) = \omega^i$ for every i . Thus we can let χ_ω denote the unique character that maps g to ω . But now if we let $\omega = e^{2\pi i/n}$, then ω is what is known as a “primitive root of unity,” which is to say that all the roots of unity are given by $1, \omega, \omega^2, \dots, \omega^{n-1}$. Notice that these roots form a multiplicative group that is isomorphic to G . It is easy to verify that the map which sends the element g^i of G to the character χ_{ω^i} of $\hat{G}$ is an isomorphism.

In the more general case, we appeal to the structure theorem for finite abelian groups, which says that any such group G can be written as a product $G_1 \times \dots \times G_k$ of cyclic groups. This means that every element g of G can be written uniquely as a product $g = g_1 g_2 \dots g_k$, where each g_i is in G_i . Given characters $\chi_1, \chi_2, \dots, \chi_k$ on $G_1, G_2, \dots, G_k$ respectively, one gets a character χ on G defined by $\chi(g) = \chi_1(g_1) \chi_2(g_2) \dots \chi_k(g_k)$, where $g = g_1 g_2 \dots g_k$ is the decomposition described above. Moreover, it is not hard to show that every character on G arises in this way. This shows that $\hat{G}$ is isomorphic to $\hat{G}_1 \times \hat{G}_2 \times \dots \times \hat{G}_k$. By the analysis of the cyclic case, the latter is, in turn, isomorphic to $G_1 \times G_2 \times \dots \times G_k$, and hence to G .

The following theorem expresses two important properties, known as the “orthogonality relations” for group characters.

Theorem 4.4. *Let G be finite abelian group. Then for any character χ in $\hat{G}$, we have*

$$\sum_{g \in G} \chi(g) = \begin{cases} |G| & \text{if } \chi = \chi_0 \\ 0 & \text{if } \chi \neq \chi_0, \end{cases}$$

and for any element g of G , we have

$$\sum_{\chi \in \hat{G}} \chi(g) = \begin{cases} |G| & \text{if } g = 1_G \\ 0 & \text{if } g \neq 1_G. \end{cases}$$

The first equation clearly holds when χ is the trivial character, χ_0 , since, in this case, each term of the sum is equal to 1. Otherwise, pick h such that $\chi(h) \neq 1$ and note

$$\chi(h) \sum_{g \in G} \chi(g) = \sum_{g \in G} \chi(hg) = \sum_{g \in G} \chi(g).$$

Since $\chi(h) \neq 1$, we must have $\sum_{g \in G} \chi(g) = 0$. The second equation can be established in a similar way.

The orthogonality relations make it possible to do “finite Fourier analysis,” in the following sense: if f is any function from G to the complex numbers and we define the “Fourier transform” $\hat{f}$ of f by $\hat{f}(\chi) = \sum_g f(g) \chi(g)$, then f can be recovered from its Fourier transform: $f = \frac{1}{|G|} \sum_\chi \hat{f}(\chi) \chi$. This shows, in particular, that any function from G to the complex numbers can be written

as a linear combination of characters. The second orthogonality relation also provides the following useful corollary:

Corollary 4.5. *For any $g, h \in G$ we have the following:*

$$\sum_{\chi \in \widehat{G}} \chi(g) \overline{\chi(h)} = \begin{cases} |G| & \text{if } g = h \\ 0 & \text{if } g \neq h \end{cases}$$

This follows from the fact that we have

$$\sum_{\chi \in \widehat{G}} \chi(g) \overline{\chi(h)} = \sum_{\chi \in \widehat{G}} \chi(g) \chi(h)^{-1} = \sum_{\chi \in \widehat{G}} \chi(gh^{-1}) = \begin{cases} |G| & \text{if } g = h \\ 0 & \text{if } g \neq h \end{cases}.$$

This corollary will enable us to focus on the residue class of m modulo k in the proof of Dirichlet's theorem.

4.3 Dirichlet characters and L -series

Let k be an integer greater than or equal to 1. It is a fundamental theorem of number theory that an integer n is relatively prime to k if and only if n has a multiplicative inverse modulo k ; in other words, if and only if there is some n' such that $nn' \equiv 1 \pmod{k}$. This implies that the residue classes of integers modulo k that are relatively prime to k form a group, denoted $(\mathbb{Z}/k\mathbb{Z})^*$, with multiplication modulo k . The cardinality of $(\mathbb{Z}/k\mathbb{Z})^*$, that is, the number of residues relatively prime to k , is denoted $\varphi(k)$, and the function φ is called the *Euler φ -function*.

Any character χ on $(\mathbb{Z}/k\mathbb{Z})^*$ can be “lifted” to a function X from $\mathbb{Z}$ to $\mathbb{C}$ defined by

$$X(n) = \begin{cases} \chi(n \bmod k) & \text{if } n \text{ is relatively prime to } k \\ 0 & \text{otherwise.} \end{cases}$$

Such a function is called a *Dirichlet character modulo k* . Dirichlet characters are *completely multiplicative*, which is to say, $X(mn) = X(m)X(n)$ for every m and n in $\mathbb{Z}$. Mathematicians typically use the symbol χ to range over Dirichlet characters, blurring the distinction between such functions and their group-character counterparts. This is harmless, since there is a one-to-one correspondence between the two, and so we will adopt this practice as well.

We can now generalize the method of Euler's proof. Roughly speaking, we need a variant of the zeta function that will allow us to focus on primes in a particular residue class modulo k . To that end, given a Dirichlet χ character modulo k , define the *Dirichlet L -function*, or *L -series*,

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s},$$

where s is any complex number. This formal series will converge whenever $\Re(s) > 1$, that is, the real part of s is greater than one.²⁴ And just as $\sum_{n=1}^{\infty} \frac{1}{q^s}$ can be written as a product via the Euler product formula, so each $L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}$ has a useful product expansion.

Theorem 4.6. *Let χ be a Dirichlet character modulo k . Then the L -function associated with χ has an Euler product expansion for $\Re(s) > 1$,*

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} = \prod_q \left(1 - \frac{\chi(q)}{q^s}\right)^{-1} = \prod_{q \nmid k} \left(1 - \frac{\chi(q)}{q^s}\right)^{-1}.$$

The last identity follows from the fact that, in the product, we can ignore those primes q that divide k , since $\chi(q) = 0$ for such q . With the product formula in place, we can sketch a proof of Dirichlet's theorem.

4.4 Contemporary proofs of Dirichlet's theorem

Recall that we want to prove that there are infinitely many primes q such that $q \equiv m \pmod{k}$, where m and k are relatively prime. As in the proof that $\sum_q \frac{1}{q^s}$ diverges, we begin by taking logarithms of both sides of the Euler product expansion for $L(s, \chi)$, where χ is a Dirichlet character modulo q :

$$\log L(s, \chi) = - \sum_{q \nmid k} \log \left(1 - \frac{\chi(q)}{q^s}\right).$$

As before, we make use of the Taylor series expansion for the logarithm on the right hand side of the above equation to obtain:

$$\begin{aligned} \log L(s, \chi) &= \sum_{q \nmid k} \sum_{j=1}^{\infty} \frac{1}{j} \frac{\chi(q^j)}{q^{sj}} \\ &= \sum_{q \nmid k} \frac{\chi(q)}{q^s} + \sum_{q \nmid k, j=2}^{\infty} \frac{1}{j} \frac{\chi(q^j)}{q^{sj}}. \end{aligned}$$

One can show that the second term in the expression is bounded by a constant that is independent of s and χ , which can be expressed as follows:

$$\log L(s, \chi) = \sum_{q \nmid k} \frac{\chi(q)}{q^s} + O(1).$$

Now comes the crucial use of Corollary 4.5 to pick out the primes in the relevant residue class. We multiply each side of the above equation by $\overline{\chi(m)}$ and then take the sum of these over all the Dirichlet characters modulo k . (Recall that we

²⁴By analytic continuation, each of these functions except for $L(s, \chi_0)$ can be extended to an analytic function on the domain $\{s \mid \Re(s) > 0\}$.

can identify each character Dirichlet with the corresponding group character, that is, the corresponding element of $(\mathbb{Z}/k\mathbb{Z})^*$.) Thus we have:

$$\sum_{\chi \in (\mathbb{Z}/k\mathbb{Z})^*} \overline{\chi(m)} \log L(s, \chi) = \sum_{\chi \in (\mathbb{Z}/k\mathbb{Z})^*} \overline{\chi(m)} \sum_{q \nmid k} \frac{\chi(q)}{q^s} + O(1).$$

To simplify this expression, we exchange the summations on the right-hand side, and appeal to Corollary 4.5. Since the cardinality of the group $(\mathbb{Z}/k\mathbb{Z})^*$ is $\varphi(k)$, we obtain

$$\sum_{\chi \in (\mathbb{Z}/k\mathbb{Z})^*} \overline{\chi(m)} \log L(s, \chi) = \varphi(k) \sum_{\substack{q \equiv m \\ (\text{mod } k)}} \frac{1}{q^s} + O(1). \quad (2)$$

This is analogous to the equation (1) in Euler's proof. Our goal is once again to show that the left-hand side tends to infinity as s approaches 1 from above; this implies that the right-hand side tend to infinity, which, in turn, implies that there infinitely many primes q that are congruent to m modulo k . However, now the left-hand side is considerably more complicated than the expression $\log \sum_{n=1}^{\infty} \frac{1}{n^s}$ in Euler's proof.

To show that $\sum_{\chi \in (\mathbb{Z}/k\mathbb{Z})^*} \overline{\chi(m)} \log L(s, \chi)$ tends to infinity as s approaches 1, we divide the characters into three classes, as follows:

1. The first class contains only the principal character χ_0 , which takes the value of 1 for all arguments that are relatively prime to k , and 0 otherwise.
2. The second class consists of all those characters which take only real values (i.e. 0 or ± 1), other than the principal character.
3. The third class consists of those characters which take at least one complex value.

It is not difficult to show that $L(s, \chi_0)$ has a simple pole at $s = 1$, which implies that the term $\overline{\chi_0(m)} \log L(s, \chi_0)$ approaches infinity as s approaches 1. The real work involves showing that for all the other characters χ , $L(s, \chi)$ has a finite nonzero limit. This implies that the other terms in the sum approach a finite limit, and so the entire sum approaches infinity.

For characters in the third class, that is, the characters that take on at least one complex value, the result is not difficult. For characters in the second class, the result is much harder, and Dirichlet used deep techniques from the theory of quadratic forms to obtain it. In the years that followed, other mathematicians found alternative, and simpler, ways of handling this case. But even in modern presentations, this case remains the most substantial and technically involved part of the proof.

Our presentation has been thoroughly "modern." In the next section, we will consider some of the methodological features of the proof that make it so. This will enable us to draw interesting contrasts with Dirichlet's proof, and then explore the way that presentations of Dirichlet's theorem gradually took on such a modern character.

5 Modern aspects of contemporary proofs

5.1 The hermeneutics of proof

Although our presentation uses contemporary terminology and notation, there is a sense in which it is a faithful description of Dirichlet’s 1837 proof. Dirichlet did not, and could not, rely on a general notion of group character, as the general notion of a group was not articulated before Cayley did so in 1854 [15], and was not brought into general currency until Kronecker’s 1870 paper [63], which first presented the structure theorem for finite abelian groups.²⁵ Another important difference is that even though Dirichlet’s argument used complex numbers in a central way, there was less established background in complex analysis than is available today, and Dirichlet tended to reduce the calculations to real analysis whenever possible. Thus his variable s ranged over real numbers, and his calculations involve real-valued sines and cosines where today we are comfortable sticking with the complex exponential. In addition, we have already noted that many of the technical details were streamlined over the years. Despite all this, the outline above characterizes the central ideas of his proof, and most mathematicians would not find it unreasonable to say that that is, essentially, how he obtained the result.

But, as we will see in Section 6, there is one very striking difference: in Dirichlet’s original presentations, there is no mention of characters at all. That is, Dirichlet’s papers contain certain expressions that we now recognize as values of the various characters, and summations that are *tantamount* to summing over all the characters. But the characters themselves are only objects that we project back into the argument from our current understanding. They hover over the page as shade-like premonitions, ghosts of mathematics yet to come.

In Section 7, we will consider presentations of Dirichlet’s theorem given by Dedekind, de la Vallée-Poussin, Hadamard, Kronecker, and Landau, and see how the characters were gradually brought to life. We will see that many of the benefits of giving characters a substantive embodiment are notational and pragmatic, but that is not to say that they are *merely* notational and pragmatic: treating characters as bona fide objects comes with serious mathematical constraints and obligations, and provides conceptual reorientations that have great bearing on the kind of mathematics we do, and the way we do it. We will argue that the reification of the notion of a character is a prototypical instance of the conceptual changes that are hallmarks of the transition to “modern” mathematical thought, and that understanding how and why the changes came about shed light on the way we do mathematics today.

But the observations above present us with a terminological conundrum: should we describe the various historical texts as “versions” or “presentations” of Dirichlet’s proof, or different proofs entirely? Having raised this issue, we will, for the most part, set it aside, and be fairly cavalier with our terminology. Since our specific concern is to study the way that language, conceptualization, and inferential practice evolved over the years, and the effects that had on the

²⁵See Wussing [106] for the history of group theory.

mathematics, we need not explicitly address the question as to when it is proper to consider two proofs essentially the same or essentially different.

Talking about historical texts in modern terms is difficult, and it is always misleading to portray the history of mathematics as a muddled and inefficient attempt to arrive at the contemporary enlightened view. We hope we have not fallen into this trap. If there is anything that deserves to be treated as a rational pursuit, mathematics should count as such, and so one would expect there to be good reasons that we do mathematics the way we do. At the same time, there are often good reasons to question the way we do mathematics today, and recognize there are tradeoffs involved in the historical decisions that were made. Comparing mathematical texts from different historical eras and trying to understand what has changed provides with a fruitful way of understanding the values that drive mathematical change. But it is often easier to draw contrasts by starting with the mathematics with which we are most familiar, and so, having discussed contemporary approaches to Dirichlet's theorem, let us foreshadow some of the contrasts we wish to consider.

5.2 Characters as objects

In Section 3, we noted that characters (whether we refer to group characters, or Dirichlet characters) are instances of the contemporary function concept, a concept which evolved significantly over the course of the nineteenth century. These are some of the salient features of the treatment of characters in our modern presentation:

1. Group characters are given an abstract, axiomatic definition as functions that satisfy the homomorphism property, and the Dirichlet characters are introduced as a natural extension of the notion.
2. In particular, one defines the set of characters modulo k extensionally. Only later does one show that this set is finite, and provide explicit ways of describing and enumerating them.
3. Characters are studied in their own right, and their general properties are enunciated in propositions and theorems.
4. One sums over sets of characters, without needing representations for any particular one. More, generally, one characterizes operations on characters (such as the product of two characters) extensionally, and not in terms of their representations.
5. Characters appear as arguments to other functions, namely, the L -functions. In particular, it is clear from the definition that $L(s, \chi)$ depends only on the extension of χ .
6. One defines various sets of characters extensionally, for example, distinguishing the trivial, real, and complex characters in terms of the values they assume. More generally, one typically carries out arguments without making reference to any particular representation.

7. The characters modulo k are viewed as elements of an algebraic structure, namely, a group, with multiplication defined pointwise and the trivial character serving as identity.

In Section 6, we will discuss Dirichlet’s original proof, and in Section 7, we will consider the way the proof was gradually transformed to reflect our contemporary understanding. We will see that proofs along the way possess various subsets of the properties just enumerated, and that, in some cases, the authors are noticeably squeamish, or at least self-conscious, of these features. Importantly, Dirichlet’s original proof has *none* of the properties just enumerated, providing a clear contrast to the style of presentation that is common today.

One way of characterizing the difference between Dirichlet’s original proof and our modern presentation is to say that, in the latter, characters are treated as full-fledged mathematical objects, whereas there are no such objects in Dirichlet’s version. The claim that over the course of the century characters gradually become treated as new sorts of objects supports our contention that the transformation has metaphysical overtones. But what, exactly, does it mean to say that a piece of mathematics sanctions certain entities as “objects”? We do not claim to have a precise set of criteria, but the list above provides some insight as to potential hallmarks of object-hood.

To start with, consider the fact that in the modern proof we identify the concept of a “character,” reason about entities falling under this concept, and ascribe various properties to them. This seems to be a bare-minimum requirement to support the claim that a mathematical text sanctions certain entities as objects, namely, that it recognizes them as being entities of a certain *sort*, capable of bearing predicates and being the target of certain operations. It doesn’t matter whether we take this sort as fundamental (for example, as we take the notion of “integer” in most contexts) or as derived from a broader sort (for example, when we view characters as functions of a certain kind). What is important is that the entities belong to a grammatically recognized category, and this category helps determine the predicates and operations that can be meaningfully ascribed to it. For example, one can talk about one integer being divisible by another, but not one character as being divisible by another. In sum, the first criterion to look for is whether the entities in question have a recognizable role in the grammar of the language.

The use of the word “representation” in our informal list provides another clue, insofar as we generally speak of a representation *of* something or other. We think of expressions like “6” and “ 2×3 ” as representing an integer. As Michael Detlefsen has pointed out to us, one common view is that an “object” is what remains invariant under all the different representations of a thing; in other words, what is left over when one has “squeezed out” all the features that are contingent on particular representations. When it comes to the notion of a function, what is the underlying invariant? There may be lots of ways of representing a particular function, but what makes them representations of the *same* function is surely that they take the same values on any given input. Thus reasoning about functions extensionally is a sign that one is reasoning

about functions *as* objects, rather than reasoning about their representations.²⁶

A third hallmark of object-hood that is present in our list is evidenced by the fact that we can *sum* over characters, just as we can sum over natural numbers. Notice that in an expression $\sum_{\chi} \dots \chi \dots$, the variable χ is a bound variable that ranges over the entities in question. Similar considerations hold for the universal and existential quantifiers. Viewing the natural numbers as quintessential mathematical objects, a sign that an entity has attained the status of object-hood is that it is possible to quantify over them in theorems and definitions, just as one quantifies over the natural numbers.²⁷ The consideration admits of degrees: whereas the bare-minimum requirement discussed above may allow us to state theorems about, and define operations on, “arbitrary” entities of the sort, a more full-blown notion of object-hood will give us more latitude in the kinds of quantification and binding that are allowed.

A fourth criterion for object-hood is evidenced by the fact that characters are allowed to appear as *arguments* to the L -functions, for example, in the expression $L(s, \chi)$. To avoid making this consideration depend on the modern notion of a function, let us note that what is essential here that an expression denoting a recognized mathematical object (in this case, a complex number) is allowed to *depend* on a character, much the way that a real number $(s)_i$ in a sequence depends on the value of the index i , or, in Euler’s notation, the value πn depends on n . What makes this more potent than the mere ability to define operations on characters is that the dependent expressions are treated as objects in their own right. $L(s, \chi)$ is not just an operation on s and χ : fixing χ , the function $s \mapsto L(s, \chi)$ is an object that one can integrate, and summing over χ , one obtains a function of s . In a similar way, one can form sets and sequences of characters, in much the same way that one forms sets and sequences of numbers, and one can define a group whose elements are characters, in much that same way that one can form a group whose elements are residues modulo some number m .

To summarize, here are some of the various senses in which characters are treated as objects in the modern presentation:

1. Characters fall under a recognized grammatic category, which allows us to state things about them and define operations and predicates on them.
2. There is a clear understanding of what it means for two expressions to represent the *same* character, and conventions ensure that the expressions occurring in a proof respect this “sameness.”
3. One can quantify and sum over characters; in short, they can fall under the range of a bound variable.
4. One can define functions which take characters as arguments, and sets of characters; indeed, characters can be elements of arbitrary algebraic structures.

²⁶Recall Quine’s dictum that “there is no entity without identity,” for example in [87].

²⁷This echoes another Quine dictum, “to be is to be the value of a bound variable” [89, p. 15].

We recognize that determining the “ontological commitments” of a practice may not be as clear-cut as Quine’s writings suggest, and we do not claim to have given a precise sense to the question as to whether a particular mathematical proof is committed to functions as objects. But we do claim to have identified various important senses in which contemporary proofs of Dirichlet’s theorem treat functions as ordinary mathematical objects, whereas Dirichlet’s original proof does not.

Below, we will not directly address the question as to whether any of these are necessary or sufficient criteria for being a mathematical object. After all, our goal is not so much to clarify the notion of a mathematical object as to explore the reasons that mathematical language and practice evolved the way it did, and the rational considerations that bore upon this evolution. But we hope we have made clear that these features of the treatment of characters turn on very fundamental aspects of mathematical language and inferential practice, and, as such, reflect nontrivial metaphysical commitments.

6 Dirichlet’s original proof

We have already asserted that characters appear only “implicitly” in Dirichlet’s original proof [28]. There is nothing mysterious about this: what we mean is that, in Dirichlet’s proof, there are certain symbolic expressions that we now recognize as denoting the values of characters; and that, moreover, some of Dirichlet’s calculations and inferences invoke what we now recognize as general properties of characters.

Let us spell out the details. Like Dirichlet, we will first consider the case where the common difference in the arithmetic progression is a prime number, denoted by p instead of k . It was well known in the nineteenth century that one can always find a *primitive element* modulo p , which is to say, an element c , such that the $p - 1$ residues $c^0, c^1, c^2, \dots, c^{p-2}$ modulo p yields all the nonzero residues, $1, 2, \dots, p - 1$ (not necessarily in the same order). In modern terms, we would say that the group $(\mathbb{Z}/p\mathbb{Z})^*$ of units modulo p is cyclic, generated by the residue class of c . In more elementary terms, this amounts to saying that for every integer n , there is a number γ_n with the property that $c^{\gamma_n} \equiv n \pmod{p}$. We saw in Section 4.2 that if χ is a Dirichlet character modulo p (that is, χ corresponds to character on $(\mathbb{Z}/p\mathbb{Z})^*$), then $\chi(c)$ is a p th root of unity, say, ω ; and, moreover, χ is entirely determined by ω , in the sense that for every n relatively prime to p , $\chi(n) = \omega^{\gamma_n}$. Thus Dirichlet simply wrote ω^{γ_n} where we would write $\chi(n)$. The notation presupposes that one has fixed a choice of the primitive element, c , though any primitive element will work equally well.

So far, so good. In the more general case where the modulus is a composite number k , however, things get more complicated. First, write k as a product of primes,

$$k = 2^\lambda p_1^{\pi_1} p_2^{\pi_2} \cdots p_j^{\pi_j}$$

where each p_i is an odd prime and π_i is greater than or equal to 1. Then the group of units modulo k is isomorphic to the product of the groups of units

modulo each term in the factor. Gauss had already shown if p is an odd prime and π is an integer greater than or equal to 1, then one can more generally find a primitive element c modulo p^π . This means that the residue class of c generates the cyclic group $(\mathbb{Z}/p^\pi\mathbb{Z})^*$, or, equivalently, for every n relatively prime to p there is a γ_n such that $c^{\gamma_n} \equiv n \pmod{p^\pi}$. Thus we can choose primitive elements $c_1, \dots, c_j$ corresponding to $p_1^{\pi_1}, p_2^{\pi_2}, \dots, p_j^{\pi_j}$. If $\lambda \geq 3$, however, there is no primitive element modulo 2^λ . Rather, $(\mathbb{Z}/2^\lambda)^*$ is a product of two cyclic groups, and for every n relatively prime to 2^λ there are an α_n and β_n such that $(-1)^{\alpha_n} 5^{\beta_n} \equiv n \pmod{2^\lambda}$. Thus for any n relatively prime to k , we can write

$$n \equiv (-1)^{\alpha_n} 5^{\beta_n} c_1^{\gamma_{1,n}} c_2^{\gamma_{2,n}} \dots c_j^{\gamma_{j,n}} \pmod{k}$$

where each $\gamma_{i,n}$ is the index n relative to $p_i^{\pi_i}$. As above, if we choose appropriate roots of unity $\theta, \varphi, \omega_1, \omega_2, \dots, \omega_j$, we obtain a character

$$\chi(n) = \theta^{\alpha_n} \varphi^{\beta_n} \omega_1^{\gamma_{1,n}} \omega_2^{\gamma_{2,n}} \dots \omega_j^{\gamma_{j,n}}. \quad (3)$$

And, once again, every character is obtained in this way. We should note that Dirichlet used the notation $p, p', \dots$ rather than $p_1, \dots, p_j$ to denote the sequence of odd primes. Moreover, he used the notation $\alpha, \beta, \gamma, \gamma', \dots$ to denote the indices, suppressing the dependence on n . Thus, Dirichlet wrote $\theta^\alpha \varphi^\beta \omega^\gamma \omega'^{\gamma'} \dots$ for the expression we have denoted $\chi(n)$ above, leaving it up to us to keep in mind that $\alpha, \beta, \dots$ depend on n .

To summarize, in the simple case of a prime modulus p , Dirichlet fixed a primitive element modulo c , and represented each character χ in terms of a p th root of unity, ω . In that case, the value $\chi(n)$ is given by ω^{γ_n} . In the more general case of a composite modulus k , Dirichlet fixed primitive elements modulo the terms of the prime factorization of k , and represented each character χ in terms of a sequence $\theta, \varphi, \pi, \pi'$ of roots of unity. In that case, the value $\chi(n)$ was written $\theta^\alpha \varphi^\beta \omega^\gamma \omega'^{\gamma'} \dots$, suppressing the information that the exponents $\alpha, \beta, \gamma, \gamma', \dots$ depend on n .

Recall that our contemporary presentation had little to say about *particular* characters, other than the trivial character, χ_0 . Rather, characters appear as arguments to the L -functions, $L(s, \chi)$, and the proof has us consider summations over the set of all characters. Let us now consider how Dirichlet handled these as well.

Again, with Dirichlet, we begin with the easier case where the common difference of the arithmetic progression is a prime, p . Recall that, in that case, each character χ corresponds to a p th root of unity, ω . Dirichlet stated the Euler product formula as follows:

We therefore have the equation

$$\prod \frac{1}{1 - \omega^\gamma \frac{1}{q^s}} = \sum \omega^\gamma \frac{1}{n^s} = L, \quad (4)$$

where the multiplication sign ranges over the whole series of primes with the sole exception of p , while the summation involves all the

integers from 1 to ∞ that are not divisible by p . The letter γ denotes γ_p on the left, and γ_n on the right.²⁸ [28, p. 3]

Compare this to the statement of Theorem 4.6 above. Since there are $p - 1$ distinct $p - 1^{st}$ roots of unity, Dirichlet continued:

The equation just found represents $p - 1$ different equations, which are obtained by replacing ω with its $p - 1$ values. It is known that these $p - 1$ different values can be represented as powers of one such Ω , chosen appropriately, so that the values are then:

$$\Omega^0, \Omega^1, \Omega^2, \dots, \Omega^{p-2}.$$

In accordance with this representation, we will write the different values L of the series or product as:

$$L_0, L_1, L_2, \dots, L_{p-2},$$

...²⁹ [28, p. 3]

Notice that Dirichlet says that the Euler product formula “represents $p - 1$ different equations,” rather than thinking of it as a single equation parametrized by ω .

In the more general case where the common difference is some composite k , Dirichlet’s procedure is completely analogous. First, he demonstrated that the Euler product formula holds:

$$\prod \frac{1}{1 - \theta^\alpha \varphi^\beta \omega^\gamma \omega'^{\gamma'} \dots \frac{1}{q^s}} = \sum \theta^\alpha \varphi^\beta \omega^\gamma \omega'^{\gamma'} \dots \frac{1}{n^s} = L, \quad (5)$$

²⁸“Man hat daher die Gleichung:

$$\prod \frac{1}{1 - \omega^\gamma \frac{1}{q^s}} = \sum \omega^\gamma \frac{1}{n^s} = L, \quad (4)$$

wo sich die Multiplicationszeichen auf die ganze Reihe der Primzahlen, mit alleiniger Ausnahme von p , erstreckt, während die Summation sich auf alle ganzen Zahlen von 1 bis ∞ bezieht, welche nicht durch p teilbar sind. Der Buchstabe γ bedeutet auf der ersten Seite γ_q , auf der zweiten dagegen γ_n .” [28, pp. 317–318] We have replaced Dirichlet’s equation number with our own, and throughout this section we have modified the translation cited in [28].

²⁹“Die eben gefundene Gleichung repräsentirt $p - 1$ verschiedene Gleichungen, welche man erhält, wenn man für ω seine $p - 1$ Werthe setzt. Bekanntlich lassen sich diese $p - 1$ verschiedenen Werthe durch die Potenzen von einem derselben Ω darstellen, wenn dieser gehörig gewählt wird, und sind dann:

$$\Omega^0, \Omega^1, \Omega^2, \dots, \Omega^{p-2}.$$

Wir werden, dieser Darstellung entsprechend, die verschiedenen Werthe L der Reihe oder des Productes mit:

$$L_0, L_1, L_2, \dots, L_{p-2}$$

bezeichnen...” [28, p. 318]

where the multiplication sign ranges over all primes, with the exclusion of $2, p, p', \dots$, and the summation ranges over all the positive integers that are not divisible by any of the primes $2, p, p', \dots$. The system of indices $\alpha, \beta, \gamma, \gamma', \dots$ on the left side corresponds to the number q , and on the right side to the number n . The general equation (5), in which the different roots $\theta, \phi, \omega, \omega', \dots$ can be combined with one another arbitrarily, clearly contains K -many particular equations.³⁰ [28, p. 17]

Note, again, Dirichlet's characterization of the general equation as "containing" the particular instances. Here, K is what we have called $\varphi(k)$, the cardinality of the group $(\mathbb{Z}/k\mathbb{Z})^*$. Dirichlet went on to note that we can choose primitive roots of unity $\Theta, \Phi, \Omega, \Omega', \dots$ so that all choices of $\theta, \varphi, \omega, \omega', \dots$ can be expressed as powers of these,

$$\theta = \Theta^a, \varphi = \Phi^b, \omega = \Omega^c, \omega' = \Omega^{c'}, \dots,$$

just as in the simpler case. He wrote that we can thus refer to the L -series in a "convenient" (*bequem*) way, as $L_{a,b,c,c',\dots}$, where $a, b, c, c', \dots$ are the exponents of the chosen primitive roots. Notice that the representations just described depend on fixed, but arbitrary, choices of the primitive roots of unity, as well as fixed but arbitrary generators of the cyclic groups. Modulo those choices, we have parameters $a, b, c, c', \dots$ that vary to give us all the characters; and for each choice of $a, b, c, c', \dots$ we have an explicit expression that tells us the value of the character at n . For Dirichlet, summing over characters therefore amounted to summing over all possible choices of this representing data.

In the special case of where the common difference is a prime, p , Dirichlet ran through calculations similar to those described in Section 4.4 to obtain the following identity:

$$\begin{aligned} & \sum \frac{1}{q^{1+p}} + \frac{1}{2} \sum \frac{1}{q^{2+2p}} + \frac{1}{3} \sum \frac{1}{q^{3+3p}} + \dots \\ &= \frac{1}{p-1} (\log L_0 + \Omega^{-\gamma_m} \log L_1 + \Omega^{-2\gamma_m} \log L_2 + \dots + \Omega^{-(p-1)\gamma_m} \log L_{p-2}). \end{aligned}$$

He has essentially arrived at equation (2) in Section 4.4, which read as follows:

$$\sum_{\chi \in (\mathbb{Z}/k\mathbb{Z})^*} \overline{\chi(m)} \log L(s, \chi) = \varphi(k) \sum_{\substack{q \equiv m \\ (\text{mod } k)}} \frac{1}{q^s} + O(1) \quad (2)$$

³⁰ “

$$\prod \frac{1}{1 - \theta^\alpha \varphi^\beta \omega^\gamma \omega'^{\gamma'} \dots \frac{1}{q^s}} = \sum \theta^\alpha \varphi^\beta \omega^\gamma \omega'^{\gamma'} \dots \frac{1}{n^s} = L, \quad (5)$$

wo sich das Multiplicationszeichen auf die ganze Reihe der Primzahlen, mit Ausschluss von $2, p, p', \dots$, und das Summenzeichen auf alle positiven ganzen Zahlen, welche durch keine der Primzahlen $2, p, p', \dots$ theilbar sind, erstreckt. Das System der Indices $\alpha, \beta, \gamma, \gamma', \dots$ entspricht auf der ersten Seite der Zahl q , auf der zweiten Seite der Zahl n . Die allgemeine Gleichung (5), in welcher die verschiedenen Wurzeln $\theta, \phi, \omega, \omega', \dots$ auf irgend eine Weise mit einander combinirt werden können, enthält offenbar eine Anzahl K besonderer Gleichungen.” [28, pp. 336–337] We have replaced Dirichlet's equation number with our own.

In the case at hand, k is p , in which case $\varphi(k)$, the cardinality of $(\mathbb{Z}/k\mathbb{Z})^*$, is equal to $p - 1$. To facilitate the comparison, switch the right-and-side with the left-hand-side of (2) (convention dictates that the term $O(1)$ stays on the right side), divide through by $\varphi(k)$, and note all of the following. First, Dirichlet used $1 + \rho$ in place of s to denote the quantity that approaches 1 from above. Second, the sum

$$\frac{1}{2} \sum \frac{1}{q^{2+2\rho}} + \frac{1}{3} \sum \frac{1}{q^{3+3\rho}} + \dots$$

has been absorbed into the term $O(1)$ in equation (2); indeed, Dirichlet's next move was to note that this sum is bounded by a constant. Finally, if $\chi_i(m)$ is equal to $\Omega^{i\gamma_m}$, then the complex conjugate $\overline{\chi_i(m)}$ is equal to $\Omega^{-i\gamma_m}$, so the expression $\Omega^{-i\gamma_m} \log L_i$ would be expressed in our notation as $\overline{\chi_i(m)} L(s, \chi_i)$.

When Dirichlet considered the more general case, he arrived at the analogous result:

$$\begin{aligned} \sum \frac{1}{q^{1+\rho}} + \frac{1}{2} \sum \frac{1}{q^{2+2\rho}} + \frac{1}{3} \sum \frac{1}{q^{3+3\rho}} + \dots \\ = \frac{1}{K} \sum \Theta^{-\alpha_m \mathfrak{a}} \Phi^{-\beta_m \mathfrak{b}} \Omega^{-\gamma_m \mathfrak{c}} \Omega^{-\gamma'_m \mathfrak{c}'} \dots \log L_{\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{c}', \dots} \end{aligned}$$

Here the summation on the right hand side of the equation is over the possible values of $\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{c}', \dots$. Once again, this translates to our equation (2).

Let us now summarize the differences between Dirichlet's presentation and contemporary ones. One salient difference is at the level of algebraic abstraction. The contemporary presentation developed the general notion of a character of an arbitrary finite abelian group, G , and showed that the characters themselves form a group, $\hat{G}$, isomorphic to G itself. Dirichlet, in contrast, focused on a very particular group, $(\mathbb{Z}/k\mathbb{Z})^*$, the multiplicative group of units modulo k . His presentation showed an intimate familiarity with the structure of that group, and the explicit mapping from that group to the group of characters. These details are inessential in the modern presentation.

But there is another abstraction that distinguishes modern presentations from Dirichlet's, namely, the willingness to treat characters as mathematical objects in their own right. This is, in part, facilitated by the algebraic abstraction: it is easier and more advantageous to treat characters as objects that transcend their representations when there are language and methods available that obviate the need to return to particular representations whenever there is real work to be done. But the dependence goes both ways: the abstract algebraic methods cannot even be invoked until one is willing to consider characters as objects that can bear properties and algebraic structure.

It is the absence of these two forms of abstraction that puts Dirichlet's presentation in stark contrast with the modern one. Dirichlet did not define the notion of character at all, let alone general operations on characters; nor did he identify any of their general properties. Rather, characters only come into play as symbolic expressions in the construction of the L -functions, and their properties are derived in an ad-hoc way as the proof proceeds. The characters are

thus “intensional” objects: they are represented explicitly in terms of primitive roots of unity and generators of the corresponding residue group, and the only way of reasoning about them is in terms of this representing data.

In Dirichlet’s presentation, there is no quantifying over characters. Instead, one quantifies over their representations in terms of primitive roots of unity. Similarly, there is no direct notion of summing over characters: Dirichlet was happy to sum over finite sets of natural numbers and tuples of natural numbers, but where we would sum over a finite set of characters, Dirichlet instead summed over representations in terms of such tuples. It is notable that he presented the Euler product formula in (4) and (5) in terms of arbitrary primitive roots of unity, and then explained that these “represent” or “contain” more particular equations which can then be summed over to obtain the desired result.

Finally, it is worth drawing attention to one further consequence of the different treatment of characters in the two presentations, namely, the extent to which they make important dependences explicit. Because Dirichlet’s expressions depend on so much representational data, Dirichlet often suppresses details, to keep the expressions from getting unwieldy. Thus, he wrote ω^γ , suppressing the dependence of γ on n , where we would write $\chi(n)$ for the character χ corresponding to ω . This places a greater burden on the text of the proof, and the reader’s memory, to keep track of the relevant dependences, and, for example, the ranges of a summation. Moreover, the modern notation $L(s, \chi)$ makes it easy to track the dependence on the character χ , something that is lost in Dirichlet’s presentation. Thus in the modern expression $\chi(m)L(s, \chi)$ the role of χ is clear in both terms in the product; in Dirichlet’s expression, $\Omega^{-i\gamma_m} \log L_i$ the connection is buried in the definition of L_i .

7 Later presentations

In this section, we will examine proofs and discussions of Dirichlet’s theorem by Dedekind [29], de la Vallée-Poussin [18, 19], Hadamard [53], Kronecker [66], and Landau [68, 69]. We will focus on describing the similarities and differences between the presentations of characters and L -functions, rather than describing each of the presentations in full, since most share the same general structure.

7.1 Dedekind

In 1863, Dedekind published a book, *Vorlesungen über Zahlentheorie*, based on his notes taken from a course on number theory given by Dirichlet at Göttingen. After the lectures, he added nine “supplements,” or appendices, with material of his own. Supplement VI, in particular, contained a presentation of Dirichlet’s theorem. Dedekind extended the supplements in the second edition, published in 1871, to include his theory of algebraic ideals. That theory was, in turn, revised and expanded in the third and fourth editions, which appeared in 1879 and 1894, respectively. The earlier supplements, however, were not changed after the first edition.

Dedekind began his presentation with an overview of main steps of the proof. In particular, he proved the Euler product formula, and obtained the series expansions for the L -functions and their logarithms. But the overview deals with a more general class of L -functions than we have considered so far:

The general proof of ... [Dirichlet's theorem] is based on the consideration of a class of infinite series of the form

$$L = \sum \psi(n),$$

where n runs through all positive integers and the real or complex function $\psi(n)$ satisfies the condition

$$\psi(n)\psi(n') = \psi(nn'),$$

... [and] we always assume that $\psi(1) = 1$.³¹ [29, §132]

He then went on to focus his attention on Dirichlet characters. He used the same expressions as Dirichlet to represent the characters, but whereas Dirichlet expressed each root of unity ω involved in the construction as the power Ω^i of a single primitive root of unity, Dedekind did not bother with this step. He did not explicitly refer to the relevant expressions as “characters,” but he introduced the notation $\chi(n)$ to denote their values, and pointed out that χ is completely multiplicative:

The numerator [of $\psi(n)$] $\chi(n) = \theta^\alpha \eta^\beta \omega^\gamma \omega'^{\gamma'} \dots$ has the characteristic property $\chi(n)\chi(n') = \chi(nn')$...³² [29, §133, footnote]

Nonetheless, like Dirichlet, Dedekind showed a reluctance to quantify over characters directly. He introduced the L -functions as

$$\prod \frac{1}{1 - \psi(q)} = \sum \psi(n) = L,$$

specifying that the product is to be taken over all primes not dividing the common difference, and the sum is to be taken over all natural numbers relatively prime to the common difference. Dedekind went on to note

... that the series can exhibit quite different behavior, depending on the roots of unity $\theta, \eta, \omega, \omega', \dots$ appearing in the expression for $\psi(n)$.

³¹ “Der allgemeine Beweis dieses Satzes ... stützt sich auf die Betrachtung einer Classe von unendlichen Reihen von der Form

$$L = \sum \psi(n),$$

wo der Buchstabe n alle ganzen positiven Zahlen durchlaufen muss, und die reelle oder complexe Function $\psi(n)$ der Bedingung

$$\psi(n)\psi(n') = \psi(nn')$$

genügt ... so nehmen wir immer an, dass $\psi(1) = 1$ ist.”

³² “Der Zähler $\chi(n) = \theta^\alpha \eta^\beta \omega^\gamma \omega'^{\gamma'} \dots$ besitzt die charakteristischen Eigenschaften $\chi(n)\chi(n') = \chi(nn')$...”

Since these roots can have $a, b, c, c', \dots$ values, respectively, the form L contains altogether

$$abcc' \dots = \varphi(k)$$

different particular series...³³ [29, §133]

Here, the use of the word “contains” is strongly reminiscent of Dirichlet’s language.

Dedekind then proceeded to divide the L -functions into classes. Given that the characters are defined in terms of a sequence of roots of unity, there are three distinct possibilities that can obtain for a given series L :

1. The roots of unity in the construction of the character occurring in L are all 1. There is only one such L -function, which is denoted L_1 .
2. The roots of unity in the construction of the character occurring in L are all real, i.e. are all ± 1 . L -functions that fall into this category are written as L_2 .
3. At least one of the roots of unity in the construction of the character occurring in L is imaginary. L -functions that fall into this category are written as L_3 .

Moreover, each L -function that falls into the third category has a conjugate: if $L_3 = \sum \frac{\theta^\alpha \eta^\beta \omega^\gamma \omega'^{\gamma'} \dots}{n^s}$, then there is a corresponding L -series in the same class, denoted by L'_3 , such that $L'_3 = \frac{\theta^{-\alpha} \eta^{-\beta} \omega^{-\gamma} \omega'^{-\gamma'} \dots}{n^s}$.

Let us now see how Dedekind’s notation plays out when it comes to summing over the characters. When proving that the L -functions corresponding to a complex character have a finite non-zero value as s tends to infinity, he obtained the following equation:

$$\begin{aligned} \varphi(k) \left(\sum \frac{1}{q^s} + \frac{1}{2} \sum \frac{1}{q^{2s}} + \dots + \frac{1}{\mu} \sum \frac{1}{q^{\mu s}} + \dots \right) \\ = \log L_1 + \sum \log(L_2) + \sum \log(L_3 L'_3), \quad (6) \end{aligned}$$

where, on the left hand side, the successive sums are over all the primes q not dividing k which satisfy the successive conditions $q \equiv 1 \pmod{k}$, $q^2 \equiv 1 \pmod{k}$, etc. On the right hand side the first sum

³³“Wir bemerken zunächst, dass diese Reihen je nach der Wahl der in dem Ausdrucke $\psi(n)$ vorkommenden Einheits-Wurzeln $\theta, \eta, \omega, \omega', \dots$ ein ganz verschiedenes Verhalten zeigen; da diese Wurzeln resp. $a, b, c, c', \dots$ verschiedene Werthe haben können, so sind in der Form L im Ganzen

$$abcc' \dots = \varphi(k)$$

verschiedene besondere Reihen enthalten ...”

is over all series L_2 of the second class, and the second sum is over all conjugate pairs $L_3L'_3$ of series of the third class.³⁴ [29, §136]

In a sense, Dedekind took the summations to range not over the characters or their representations, but over the *L-functions* themselves. Later, when dealing with these sums, he came closer to Dirichlet's presentation, in that the indices of the summation range over the sequences of roots that determine the characters. However, at times he introduced convenient abbreviations. For example, given a particular collection of roots of unity $\theta, \eta, \omega, \omega', \dots$, Dedekind denoted $\theta^{-\alpha_1}\eta^{-\beta_1}\omega^{-\gamma_1}\omega'^{-\gamma'_1}\dots$ by χ , where $\alpha_1, \beta_1, \gamma_1, \gamma'_1$ stand for the indices of the first term of the progression, m . Note that while Dedekind used the notation $\chi(n)$ in a footnote, he explicitly called χ , as defined here, a *value*, and indeed in this case it stands for the value $\chi(m)$. Moreover, when he used the symbol in a summation, Dedekind was explicit that the summation ranges not over χ , but rather the roots of unity involved in the definition:

The summation of all products $\chi \log L$ therefore gives the result

$$\varphi(k) \left(\sum \frac{1}{q^s} + \frac{1}{2} \sum \frac{1}{q^{2s}} + \frac{1}{3} \sum \frac{1}{q^{3s}} + \dots \right) = \sum \chi \log L, \quad (7)$$

where the successive sums on the left hand side are over all primes q satisfying the successive conditions $q \equiv m, q^2 \equiv m, q^3 \equiv m \pmod{k}$ etc., while the sum on the right hand side is over all $\varphi(k)$ different root systems $\theta, \eta, \omega, \omega', \dots$ ³⁵ [29, §137]

In many ways, Dedekind did not stray far from Dirichlet's presentation. For the most part, his treatment of the characters was intensional, in the sense that the arguments rely on the particular representations of the characters. In other words, operations on the characters are described in terms of their representations, rather than their values; and, like Dirichlet, he viewed summations as ranging over these representations in equations (6) and (7). Similarly, he classified the characters depending on the roots involved in their representation,

34 “

$$\varphi(k) \left(\sum \frac{1}{q^s} + \frac{1}{2} \sum \frac{1}{q^{2s}} + \dots + \frac{1}{\mu} \sum \frac{1}{q^{\mu s}} + \dots \right) = \log L_1 + \sum \log(L_2) + \sum \log(L_3L'_3),$$

wo auf der linken Seite das erste, zweite Summenzeichen u.s.f. sich auf alle die in k nicht aufgehenden Primzahlen q bezieht, welche resp. den Bedingungen $q \equiv 1, q^2 \equiv 1 \pmod{k}$ u.s.f. Genüge leisten; auf der rechten Seite bezieht sich das erste Summenzeichen auf alle Reihen L_2 der zweiten Classe, das zweite auf alle verschiedenen Paare $L_3L'_3$ conjugirter Reihen dritter Classe.” We have added the equation numbers in this quotation and the next, for later reference.

³⁵ “Die Summation aller Producte $\chi \log L$ giebt daher das Resultat

$$\varphi(k) \left(\sum \frac{1}{q^s} + \frac{1}{2} \sum \frac{1}{q^{2s}} + \frac{1}{3} \sum \frac{1}{q^{3s}} + \dots \right) = \sum \chi \log L,$$

wo auf der linken Seite das erste, zweite, dritte Summenzeichen u.s.f. sich auf alle Primzahlen q bezieht, welche resp. den Bedingungen $q \equiv m, q^2 \equiv m, q^3 \equiv m \pmod{k}$ u.s.f. genügen, während das Summenzeichen auf der rechten Seite sich auf die sämmtlichen $\varphi(k)$ verschiedenen Wurzel-Systeme $\theta, \eta, \omega, \omega', \dots$.”

rather than their values. Nor did he take the expression L , for the L -functions, to depend on the characters themselves, but rather as expressions that behave differently “depending on the roots of unity” appearing in their construction. The L notation in particular does nothing to signal this dependence.

Nonetheless, he did take some key steps towards viewing the characters abstractly. To start with, Dedekind went out of his way to isolate the characters as independent of the L -series in which they appear, and, in particular, flagged them as entities satisfying certain key properties. And, at least at times, he characterized summations as ranging over the L -functions themselves, hinting at a new level of abstraction.

The use of the term “character” in the modern sense can be traced to the long Supplement XI in the 1879 edition of the Dirichlet-Dedekind *Vorlesungen*, which contains a presentation of Dedekind’s theory of ideals in algebraic number fields. Some background will be helpful here. Number theory has long been concerned with questions as to which numbers can be represented by a given algebraic expression in which the variables are taken to range over the integers. It is easy to characterize the numbers that can be represented by a linear form $ax + b$ in one variable, and quadratic reciprocity addresses the problem of which numbers can be represented by a quadratic form $ax^2 + bx + c$. When it comes to quadratic forms in two variables, things become more difficult. Fermat’s famous theorem that the odd prime numbers that can be represented by the form $x^2 + y^2$ are exactly the ones that are congruent to one modulo four is considered a gem of number theory. Euler proved this and other of Fermat’s claims in the eighteenth century, and Lagrange later extended the theory of binary quadratic forms considerably. A central contribution of Gauss’ *Disquisitiones Arithmeticae* is a complete classification of binary quadratic forms, whose study constitutes the bulk of that work. Gauss showed that it suffices to characterize “primitive” quadratic forms $ax^2 + 2bxy + cy^2$, where the second coefficient is even, and a , b , and c have no factor in common. (Quadratic forms are also classified as “indefinite,” “positive definite,” and “negative definite,” and in the discussion that follows it should be assumed that we are fixing our attention on one of these fixed kinds.) He called the value $D = b^2 - ac$ the *discriminant* of the form, and showed how to assign to each primitive quadratic form of discriminant D a finite list of values of the form ± 1 which, roughly speaking, characterizes its behavior. He called these values the *characters* of the form, and took the “genus” of a form to be the collection of all the primitive forms with the same discriminant and character.

Already in the first edition of the *Vorlesungen*, Dedekind used Gauss’ terminology in Supplement IV, “Genera of quadratic forms.” In 1879, however, Dedekind went further by showing that Gauss classification could be understood in terms of his theory of ideals. Specifically, he showed that there is a correspondence between genera of quadratic forms and equivalence classes of ideals in an associated quadratic extension of the rationals. Moreover, there is a group structure on the ideals, and Gauss’ characters correspond to characters on that group, in the modern sense. This explains the quotation in Section 3: Dedekind had simply adopted Gauss’ terminology to characterize the homomorphisms

from groups of ideals to the (nonzero) complex numbers more generally.

As we noted in Section 3.1, by 1882, Weber [100] had already begun using the term “character” with respect to arbitrary finite abelian groups. He defined a (finite) abelian group as consisting of h elements “of any type” (*iregend welcher Art*) satisfying the usual group laws. He then stated the now-familiar structure theorem:

In a finite abelian group of order h one can always choose elements $\Theta_1, \Theta_2, \dots, \Theta_\nu$ of order $n_1, n_2, \dots, n_\nu$ so that each element Θ of G can be expressed uniquely by the form

$$\Theta_1^{s_1} \Theta_2^{s_2} \dots \Theta_\nu^{s_\nu},$$

where $s_1, s_2, \dots, s_\nu$ are chosen from complete residue systems with respect to the moduli $n_1, n_2, \dots, n_\nu$.³⁶ [100, pp. 306–307]

Weber called the sequence of values $\Theta_1, \Theta_2, \dots, \Theta_\nu$ a *basis* of the group, and used them to define the characters as follows:

If we assign, to the ν elements $\Theta_1, \Theta_2, \dots, \Theta_\nu$ of such a basis, ν roots of unity $\omega_1, \omega_2, \dots, \omega_\nu$ of order $n_1, n_2, \dots, n_\nu$ respectively, then each element $\Theta = \Theta_1^{s_1} \Theta_2^{s_2} \dots \Theta_\nu^{s_\nu}$ of the group also corresponds to a particular h^{th} root of unity

$$\omega = \omega_1^{s_1} \omega_2^{s_2} \dots \omega_\nu^{s_\nu}.$$

We denote this root of unity by $\chi(\Theta)$ and call it the *character* of the element.³⁷ [100, p. 307]

Weber went on to point out that this gives rise to h distinct characters, and that each such character χ satisfies the equation

$$\chi(\theta)\chi(\theta') = \chi(\theta\theta').$$

Moreover, this last condition provides an exact characterization:

³⁶ “In einer Abel’schen Grupper G von Grade h kann man stets die Elemente $\Theta_1, \Theta_2, \dots, \Theta_\nu$ von den Graden $n_1, n_2, \dots, n_\nu$ so auswählen, dass in der Form

$$\Theta_1^{s_1} \Theta_2^{s_2} \dots \Theta_\nu^{s_\nu}$$

jedes Element Θ von G und jedes nur einmal enthalten ist, wenn $s_1, s_2, \dots, s_\nu$ je einem vollständigen Restsystem nach den Moduln $n_1, n_2, \dots, n_\nu$ entnommen werden.”

³⁷ “Ordnet man den ν Elementen $\Theta_1, \Theta_2, \dots, \Theta_\nu$ einer solchen Basis ν Einheitswurzeln $\omega_1, \omega_2, \dots, \omega_\nu$ von den Graden $n_1, n_2, \dots, n_\nu$, so entspricht auch jedem Element $\Theta = \Theta_1^{s_1} \Theta_2^{s_2} \dots \Theta_\nu^{s_\nu}$ der Gruppe eine bestimmte h^{te} Einheitswurzel ω nach der Vorschrift

$$\omega = \omega_1^{s_1} \omega_2^{s_2} \dots \omega_\nu^{s_\nu}.$$

Wir bezeichnen diese Einheitswurzel mit $\chi(\Theta)$, und nennen dieselbe den *Charakter* des Elementes Θ .”

If, conversely, $\chi(\theta)$ is a uniquely determined function of θ which satisfies [the equation above], then it is necessarily contained among these h characters.³⁸ [*ibid.*]

Thus Weber’s paper provides us not only with the modern notion of a character defined on an arbitrary finite abelian group, but also with the modern understanding that such characters constitute an instance of the function concept.

There are some small differences between Weber’s presentation and ours. In his presentation of the second orthogonality relation, Weber used ellipses rather than summation notation to sum over the characters:

For each element Θ :

$$\chi_1(\Theta) + \chi_2(\Theta) + \dots + \chi_h(\Theta) = 0,$$

except for the identity element Θ_0 , for which we have

$$\chi_1(\Theta_0) + \chi_2(\Theta_0) + \dots + \chi_h(\Theta_0) = h.³⁹$$

[100, p. 308]

This implicitly assumes that the h characters are enumerated $\chi_1, \dots, \chi_h$, though the enumeration is arbitrary. Moreover, Weber provided the explicit construction of the set of characters before the extensional characterization, whereas modern presentations usually provide the extensional characterization first. But these points are minor, and, otherwise, his presentation is little different from the one in Section 4.2.⁴⁰

Although Dedekind revised his theory of ideals substantially in the third and fourth editions of the *Vorlesungen*, he did not revise any of the supplements that appeared in the first edition. In particular, he did not take advantage of the opportunity to go back to introduce the modern notion of a character in his presentation of Dirichlet’s proof, although he clearly could have done so in the later editions.

7.2 de la Vallée-Poussin

More than two decades later, Charles Jean de la Vallée-Poussin gave another presentation of Dirichlet’s theorem in a 1895/6 paper entitled “Démonstration

³⁸Ist umgekehrt $\chi(\theta)$ eine durch das Element θ eindeutig bestimmte Function, welche der Bedingung ... genügt, so ist dieselbe nothwendig unter diesen h Charakteren enthalten.

³⁹ “Für jedes Element Θ ist:

$$\chi_1(\Theta) + \chi_2(\Theta) + \dots + \chi_h(\Theta) = 0,$$

ausgenommen für das Hauptelement Θ_0 , für welches

$$\chi_1(\Theta_0) + \chi_2(\Theta_0) + \dots + \chi_h(\Theta_0) = h.$$

⁴⁰ Mackey’s historical survey [74] of the history of harmonic analysis include a very helpful and informative overview of the history of character-theoretic ideas in number theory.

simplifiée du Théorème de Dirichlet sur la progression arithmétique,” and again in sections from his 1897 book *Recherches analytiques sur la théorie des nombres premiers*. He introduced the characters in much the same way that Dirichlet and Dedekind did, namely, via an explicit construction in terms of primitive roots and roots of unity. Like Dedekind and Weber, he used the symbol χ for the characters. Like Weber, he distinguished between then $\varphi(M)$ characters modulo M using subscripts, writing $\chi_1, \chi_2, \dots, \chi_{\varphi(M)}$.

de la Vallée-Poussin went on to subject the characters to a thorough study. Whereas Dirichlet and Dedekind divided the L -functions (or, perhaps more precisely, the corresponding expressions) into three categories based on their representations, de la Vallée Poussin based the categorization on the characters themselves. The following description applies to the simplest case, where the modulus M is a prime:

One calls the character that corresponds to the root $+1$ the *principal character*; it is equal to unity for all numbers n . Apart from the principal character, there is only one which is real for all numbers n : it corresponds to the root (-1) and is equal to ± 1 depending on the number n .

We call all the other characters *imaginary characters*, though they may have a real value for some particular numbers. Their modulus is always equal to unity.⁴¹ [19, p. 19]

Thus de la Vallée Poussin divided the characters into three classes:

1. the class consisting solely of the principal character;
2. the class consisting of all real characters (in this case, there is only one character which is real for all numbers, corresponding to the root -1); and
3. The class consisting of all other characters, called the imaginary characters.

Notice that his categorization includes both intensional and extensional characterizations; that is, he characterized the characters in each class both in terms of the values they take, and the roots involved in their construction.

In his study the characters in his 1896 paper, de la Vallée Poussin listed a number of “very important relations” (*relations très importantes*). The first is that $\chi(n)\chi(n') = \chi(nn')$ for every character χ , and n and n' . The second is that for a given character χ modulo M , and any n and n' , if $n \equiv n' \pmod{M}$ then $\chi(n) = \chi(n')$. The third and fourth are the first and second orthogonality

⁴¹ “On appelle *caractère principal* celui qui correspond à la racine $+1$; il est égal à l’unité pour tous les nombres n . En dehors du caractère principal, il n’y en a qu’un seul qui soit réel pour tous les nombres n : il correspond à la racine (-1) et est égal à ± 1 suivant le nombre n .

Nous donnerons à tous les autres caractères le nom de *caractère imaginaires*, quoiqu’ils puissent avoir une valeur réelle pour certains nombres particuliers. Leur module est toujours égal à l’unité.”

relations, respectively. In order to present the second orthogonality relation, de la Vallée Poussin introduced a new symbol, S , to denote summation over the characters:⁴²

Consider ... the sum extending over all the characters, that is to say over all the systems of roots

$$S_\chi \chi(n) = S_\omega \omega_1^{\nu_1} \omega_2^{\nu_2} \dots$$

... For every number n , the sum extending over the totality of characters satisfies

$$S_\chi \chi(n) = 0,$$

the only exception being the case where

$$n \equiv 1 \pmod{M},$$

because then all the indices are zero and one has

$$S_\chi \chi(n) = \varphi(M).$$

[18, pp. 14–15]

Although his notation is different from that of Dirichlet and Dedekind, his proof, like theirs, relies on an explicit calculation based on the construction of the characters, in contrast to the modern proof sketched in Section 4.2.

When introducing the L -functions in his 1896 paper, de la Vallée-Poussin simply described them in terms of their equivalent expressions as a sum and product. In 1897, however, he adopted a functional notation:

We define the function $Z(s, \chi \bmod M)$, for $\mathcal{R}(s) > 1$, by the absolutely convergent expressions:

$$Z(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} = \prod \left(1 - \frac{\chi(q)}{q^s} \right)^{-1}$$

⁴²“Considérons ... la somme étendue à tous les caractères, c’est-à-dire à tous les systèmes de racines

$$S_\chi \chi(n) = S_\omega \omega_1^{\nu_1} \omega_2^{\nu_2} \dots$$

... Pour tout nombre n , la somme étendue à la totalité des caractères

$$S_\chi \chi(n) = 0,$$

à la seule exception près du cas où

$$n \equiv 1 \pmod{M},$$

car alors tous les indicateurs sont nuls et l’on a

$$S_\chi \chi(n) = \varphi(M).”$$

where n designates successively all the integers prime to M and q all the prime numbers not dividing M .⁴³ [19, p. 56]

Aside from the choice of the letter Z instead of the letter L , we have finally arrived at the modern notation. But, like Dirichlet and Dedekind, he was reticent to quantify over the characters, using language that is eerily reminiscent of theirs. For example, in his 1897 work, he wrote:

... one finds the *fundamental equation*

$$(E) \dots - \lim_{s=1} (s-1) \frac{Z'(s, \chi)}{Z(s, \chi)} = \lim_{s=1} (s-1) \sum_q \chi(q) \frac{lq}{q^s},$$

and this equation (E) represents in reality $\varphi(M)$ distinct ones, which result from exchanging the characters amongst themselves.⁴⁴ [19, p. 65].

Nonetheless, there are a number of important respects in which de la Vallée's presentation is close to the modern one. To start with, characters are treated as an object of study in their own right, bearing their own properties and relations. Moreover, they are classified extensionally, although this classification is, at the same time, related to properties of their representations. They are also represented notationally as arguments to the L -functions, hinting that they are, in a sense, on par with natural numbers. Finally, and most interestingly, de la Vallée-Poussin introduced a summation notation with an index ranging over the characters, not their representing data. The fact that he went out of his way to use one symbol, S , for summation of characters, and the usual Σ symbol for summation over sets of natural numbers hints that he does not conceive of these as precisely the same mathematical operation, although they have similar properties.

7.3 Hadamard

For each real number x , let $\pi(x)$ denote the number of primes less than x . Around the turn of the century, both Legendre and Gauss conjectured that $\pi(x)$ is asymptotic to $x/\ln(x)$, in the sense that their ratio, $\pi(x) \ln x/x$, approaches 1

⁴³ "Nous définirons la fonction $Z(s, \chi \bmod M)$, pour $\Re(s) > 1$, par les expressions absolument convergentes

$$Z(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} = \prod \left(1 - \frac{\chi(q)}{q^s} \right)^{-1}$$

où n désigne successivement tous les nombres entiers premiers à M et q tous les nombres premiers qui ne divisent pas M ."

⁴⁴ "... on trouve l'équation fondamentale

$$(E) \dots - \lim_{s=1} (s-1) \frac{Z'(s, \chi)}{Z(s, \chi)} = \lim_{s=1} (s-1) \sum_q \chi(q) \frac{lq}{q^s},$$

et cette équation (E) en représente en réalité $\varphi(M)$ distinctes par l'échange des caractères entre eux."

as x approaches infinity. This fact, now known as the “prime number theorem” was finally proved by both de la Vallée-Poussin and Jacques Hadamard, working independently, in 1896. de la Vallée-Poussin’s proof was first published in the *Annales de la Société Scientifique de Bruxelles*, but also appeared in the 1897 book we discussed in the last section. Both Hadamard and de la Vallée-Poussin also obtained a generalization of the prime number theorem to arithmetic progressions, which says that if m is relatively prime to k , the number of primes less than x that are congruent to m modulo k is asymptotic to $(1/\varphi(k)) \cdot (x/\ln x)$. This is, of course, a generalization of Dirichlet’s theorem, since it implies that there are infinitely many primes congruent to m modulo k .

Hadamard’s 1896 paper was titled “Sur la distribution des zéros de la fonction $\zeta(s)$ et ses conséquences arithmétiques,” and provided on a number of results concerning L -functions and characters. He introduced characters in the same way as Dirichlet, Dedekind and de la Vallée-Poussin, namely, by a construction in terms of roots of unity and primitive elements. Like Weber and de la Vallée-Poussin, he distinguished between the characters modulo k notationally by making use of subscripts, writing $\psi_v(n)$ where v runs from 1 to $\varphi(k)$. He defined the L -functions as follows:

$$L_v(s) = \sum_{n=1}^{\infty} \frac{\psi_v(n)}{n^s}.$$

The notation $L_v(x)$, in contrast to de la Vallée-Poussin’s $Z(s, \chi)$, does not indicate an explicit dependence on the character, though the subscript provides a link. As did his predecessors, Hadamard classified the L -functions into three different classes, but his classification was intensional, like Dirichlet’s and Dedekind’s, referring to the roots of unity used in the construction of the characters.

When it came to summing over the characters, Hadamard, unlike Dirichlet, Dedekind, or de la Vallée-Poussin, let the index of the summation range over the *subscripts* described above.

The fundamental equation that Dirichlet used in the demonstration of his theorem is

$$\sum_v \frac{\log L_v(s)}{\psi_v(m)} = \varphi(k) \left(\sum \frac{1}{q^s} + \frac{1}{2} \sum' \frac{1}{q^{2s}} + \frac{1}{3} \sum'' \frac{1}{q^{3s}} + \dots \right),$$

where m is some integer prime to k , and where, among the signs $\sum, \sum', \sum'', \dots$, the first ranges over the prime numbers q such that $q \equiv m \pmod{k}$, the second ranges over the primes numbers q such that $q^2 \equiv m \pmod{k}$, etc.⁴⁵ [53, p. 209]

⁴⁵ “L’équation fondamentale utilisée par Dirichlet pour la démonstration de son théorème, est

$$\sum_v \frac{\log L_v(s)}{\psi_v(m)} = \varphi(k) \left(\sum \frac{1}{q^s} + \frac{1}{2} \sum' \frac{1}{q^{2s}} + \frac{1}{3} \sum'' \frac{1}{q^{3s}} + \dots \right)$$

où m est un entier quelconque premier avec k et où les signes $\sum, \sum', \sum'', \dots$ s’étendent, le premier aux nombres premiers q tels que $q \equiv m \pmod{k}$, le second aux nombres premiers q tels que $q^2 \equiv m \pmod{k}$, etc.”

Once again, this is comparable to the modern formulation in equation (2).

Hadamard's presentation falls somewhere between those of Dirichlet and Dedekind, on the one hand, and that of de la Vallée-Poussin, on the other. His treatment of characters was, for the most part, intensional. But, in contrast to Dirichlet and Dedekind, he left representational data out of the key summations, thereby eliminating unnecessary clutter. However, unlike de la Vallée-Poussin, he did not go so far to characterize these as summations over the characters themselves. Rather, he introduced natural number indices, v , labeling the characters, and then took the variable of summation to range over those.

7.4 Kronecker

Between 1863 and 1891, Leopold Kronecker lectured at the University of Berlin on a range of subjects, including number theory, algebra, and the theory of determinants. After Kronecker's death, his student, Kurt Hensel, edited the five volumes of his collected works, which were published between 1895 and 1930. Hensel also took it upon himself to work Kronecker's copious lecture notes and course material into two textbooks, *Vorlesungen über Zahlentheorie* and *Vorlesungen über die Theorie der Determinanten*, which he published in 1901 and 1903, respectively. The first of these closes with a proof of Dirichlet's theorem, which we will discuss here.

Kronecker, in fact, wrote his doctoral dissertation on algebraic number theory under Dirichlet's supervision, completing it in 1845. Kronecker insisted that mathematics should maintain a clear focus on symbolic representations and algorithms, a commitment that is evident throughout his work. Avoiding talk of "arbitrary" functions, real numbers, and so on, Kronecker focused instead on the construction of algebraic systems and explicit algorithms for calculating with these algebraic representations. For example, his article "Grundzüge einer arithmetischen Theorie der algebraischen Grössen" [64] provides means of carrying out operations on systems of algebraic integers in finite extensions of the rationals. Similarly, "Ein Fundamentalsatz der allgemeinen Arithmetik" [65] provides an explicit construction of a splitting field for any polynomial with integer coefficients, and he viewed this as filling a gap in Galois' work.⁴⁶

Kronecker's approach to number theory had a similar orientation. As Hensel put it in the introduction to *Vorlesungen über Zahlentheorie*:

He believed that one can and must in this domain formulate each definition in such a way that its applicability to a given quantity can be assessed by means of a finite number of tests. Likewise, an existence proof for a quantity is to be regarded as entirely rigorous only if it contains a method by which that quantity can actually be found. Kronecker was far from one to completely reject a definition or proof that does not meet these highest requirements, but he

⁴⁶See Edwards [31–34] for a discussion of these works, and Kronecker's mathematics more generally.

believed that there was then something missing, and he held that completing it along these lines is an important task, by which our knowledge is furthered in an important sense.⁴⁷ [66, p. vi]

Kronecker's proof of Dirichlet's theorem is as a focal point of the book, in that it brings together methods and ideas developed throughout the entire work. Here is Hensel's characterization:

[The book] closes with the proof of the famous theorem that any arithmetic sequence, whose first term and common difference are relatively prime, contains infinitely many prime numbers. But Kronecker completed Dirichlet's proof of this theorem in a significant sense, in that he proved that one can determine, for an arbitrarily large number μ , a larger number $\bar{\mu}$, so that in the interval $(\mu \cdots \bar{\mu})$ one is sure to find a prime number of the required form. This nice supplement to that famous proof is a fruit of the higher demands, mentioned above, that Kronecker placed on arithmetic proofs. And here it seems, in fact, that with this improvement of Dirichlet's theorem, nothing by way of simplicity or transparency has been lost.⁴⁸ [66, p. viii]

The interaction between analytic and number-theoretic methods is a central theme of the *Vorlesungen*, which opens with a discussion of Gauss' distinction between the fields of number theory, algebra, and analysis, and argues that they cannot be cleanly separated. For example, Kronecker praised Leibniz' characterization of π in terms of the series

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots = \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1}$$

as providing a definition of π of a "fully number-theoretic character" (*durchaus zahlentheoretischem Charakter*).

⁴⁷ "Er meinte, man könne und man müsse in diesem Gebiete eine jede Definition so fassen, daß durch eine endliche Anzahl von Versuchen geprüft werden kann, ob sie auf eine vorgelegte Größe anwendbar ist oder nicht. Ebenso wäre ein Existenzbeweis für eine Größe erst dann als völlig streng anzusehen, wenn er zugleich ein Method enthalte, durch welche die Größe, deren Existenz bewiesen werde, auch wirklich gefunden werden kann. Kronecker war weit davon entfernt, eine Definition oder einen Beweis vollständig zu verwerfen, der jenen höchsten Anforderungen nicht entsprach, aber er glaubte, daß dann eben noch etwas fehle, und er hielt eine Ergänzung nach dieser Richtung hin für eine wichtige Aufgabe, durch die unsere Erkenntnis in einem wesentlichen Punkte erweitert würde." We have modified and extended a translation due to Stein [94, p. 250].

⁴⁸ "...schließt mit dem Beweise des berühmten Satzes, daß jede arithmetische Reihe, deren Anfangsglied und Differenz teilerfremd sind, unendlich viele Primzahlen enthält; aber Kronecker vervollständigt den Dirichletschen Beweis dieses Satzes in einem wesentlichen Punkte, indem er nachweist, daß man für jede beliebige groß anzunehmende Zahl μ eine größere Zahl $\bar{\mu}$ so bestimmen kann, daß in dem Intervalle $(\mu \cdots \bar{\mu})$ sich sicher eine Primzahl der verlangten Form befindet. Dies schöne Ergänzung jenes berühmten Beweises ist eine Frucht der oben erwähnten höheren Forderungen, welche Kronecker an arithmetische Beweise stellte, und hier scheint es in der That, daß durch diese Verbesserung der Dirichletsche Beweis nichts an Einfachheit und Durchsichtigkeit verloren hat."

What these examples teach us holds, more generally, for all the definitions of analysis. These always lead back to the integers and their properties, and from that entire branch of mathematics, only the concept of a limit has so far remained foreign. Arithmetic cannot be separated from analysis, which has freed itself from its original source, geometry, and has developed independently, on free soil. Even less so, as Dirichlet has succeeded in obtaining the deepest and most beautiful results in arithmetic by combining the methods of the two disciplines.⁴⁹ [66, pp. 4–5]

This interplay comes to the fore in the proof of Dirichlet’s theorem. The text reports that Kronecker’s version of the proof, in the case where the common difference is prime, was worked out in the lectures he gave in the winter semester of 1875/1876, whereas the general case was presented in during the winter semester of 1886/1887 [66, p. 442]. Fixing a modulus m and an r relatively prime to m , recall that Kronecker aimed to provide, for a given number μ , an explicit upper bound $\bar{\mu}$ on the integers one has to consider to find a prime number greater than μ and congruent to r modulo m . When it came to the analytic part of the proof, Kronecker explained that obtaining the desired bound is reduced to obtaining a positive lower bound on a certain analytic series that arises in the proof. He wrote

For the ambiguous [real] characters, Dirichlet’s proof meets this requirement. But his methods are not sufficient to do the same for the series corresponding to the complex characters.⁵⁰ [66, p. 481]

He went on:

Generally speaking, this is a special case of the problem of finding, given a well-defined nonzero number, a bound, above which it necessarily lies. This is not as easy as it seems at first glance; indeed, in some circumstances, the problem can count among the thorniest questions known to science.⁵¹ [66, pp. 481–482]

Kronecker took the opportunity to clarify the methodological stance towards analysis that is appropriate to these issues. Even though, in his work, he avoided

⁴⁹ “Was uns diese Beispiele lehren, ist nun maßgebend für alle Definitionen der Analysis überhaupt. Dieselben führen stets auf die ganzen Zahlen und ihre Eigenschaften zurück, und es ist von dem ganzen Gebiete des letztgenannten Zweiges der Mathematik der einzige Begriff des limes oder der Grenze der Zahlentheorie bisher fremd geblieben. Gegen die Analysis also, die sich von ihrer ursprünglichen Quelle, der Geometrie, befreit und auf freiem Boden selbständig entwickelt hat, kann die Arithmetik nicht abgegrenzt werden, um so weniger, als es *Dirichlet* gelungen ist, grade die schönsten und tiefliegenden arithmetischen Resultate durch die Verbindung der Methoden beider Disciplinen zu erzielen.”

⁵⁰ “Für die ambigen Charaktere erfüllt der Beweis von Dirichlet auch diese Forderung, dagegen reichen seine Methoden nicht aus, um dasselbe auch für die Reihen zu leisten, welche den complexen Charakteren entsprechen.”

⁵¹ “Überhaupt ist das hier in einen speziellen Falle sich darbietende Problem, für eine von Null verschiedene wohldefinierte Zahlgröße eine Grenze zu finden, *über* der sie notwendig liegen muß, nicht so einfach, als es auf den ersten Blick erscheint, vielmehr kann diese Aufgabe unter Umständen eine der heikelsten Fragen sein, die die Wissenschaft kennt.”

the notion of an “arbitrary” real number, he recognized the importance of understanding particular number systems in analytic terms. For example, given a symbolic representation of a real number, one may wish to compute rational approximations. But knowing that a nonnegative real number is nonzero is not the same as having a positive, rational lower bound on its values.⁵² Kronecker mentioned the problem of bounding a nonzero determinant away from zero as an example of a problem that is generally difficult. He also noted that, given two convergent series, it can be difficult to determine which one has a greater value. These facts are now quite familiar in constructive and computable analysis.⁵³

Kronecker’s treatment of characters provides an interesting combination of Dirichlet’s approach and modern ones. Fixing a modulus m , Kronecker, like Dirichlet, provided a fully explicit description of the characters modulo m in terms of primitive elements of the powers of primes giving m and primitive roots of unity. However, with judicious choice of notation, he was at the same time able to suppress extraneous detail. For example, fixing primitive elements $\gamma, \gamma_0, \gamma_1, \dots, \gamma_g$, one can express any element r relatively prime to m in the form

$$r \equiv \gamma^\rho \gamma_0^{\rho_0} \dots \gamma_g^{\rho_g} \pmod{m}.$$

But then one can “package” the representing values $(\rho, \rho_0, \rho_1, \dots, \rho_g)$ as a tuple, the “index system of r ,” denoted $\text{Indd } r$. For each of the cyclic groups in the decomposition of the group of units modulo m , suppose we also choose corresponding roots of unity, $\omega, \omega_0, \omega_1, \dots, \omega_g$.⁵⁴

Now let r be a unit modulo m , and $\text{Indd } r = (\rho, \rho_0, \rho_1, \dots, \rho_g)$; we now assign to r the root of unity:

$$\Omega(r) = (-1)^\rho \omega_0^{\rho_0} \omega_1^{\rho_1} \dots \omega_g^{\rho_g}$$

which we call a *character* of r , since the index system $(\rho, \rho_0, \rho_1, \dots, \rho_g)$, and hence $\Omega(r)$, are uniquely determined.⁵⁵ [66, p. 444]

We obtain *all* the possible characters by fixing primitive roots of unity $\omega, \omega_0, \dots, \omega_g$, so that every tuple of roots can be represented in the form

$$\omega^k, \omega_0^{k_0}, \omega_1^{k_1} \dots, \omega_g^{k_g},$$

⁵²In modern terms, one can obtain such a lower bound by computing rational approximations until one obtains one that is sufficiently accurate to bound the number away from zero. Thus the statement “if $r \neq 0$, then $|r| > 0$ ” was accepted by the Russian school of constructive mathematics in the 1950’s and 1960’s. This implication is equivalent to “Markov’s principle,” which is, however, rejected by strict constructivists. See, for example, [96].

⁵³See, for example, Troelstra and van Dalen [96].

⁵⁴As above, the powers of two require special treatment. In the discussion, Kronecker assumes that m is divisible by 8, in which case the units modulo that power of two form a product of two cyclic groups; the values of ρ and ρ_0 are the indices in those two groups, and ω and ω_0 are the corresponding roots of unity.

⁵⁵Es sie nun r eine Einheit modulo m , und $\text{Indd } r = (\rho, \rho_0, \rho_1, \dots, \rho_g)$; ordnen wir r jetzt die Einheitswurzel:

$$\Omega(r) = (-1)^\rho \omega_0^{\rho_0} \omega_1^{\rho_1} \dots \omega_g^{\rho_g}$$

zu, so gehört zu jeder Einheit r eine und nur eine Einheitswurzel $\Omega(r)$, welche wir einen *Charakter* von r nennen wollen, denn durch r ist ja das Indexsystem $(\rho, \rho_0, \dots)$, also $\Omega(r)$ eindeutig bestimmt.

where the k and k_i 's are less than the cardinality of the corresponding cyclic group.

When there is no fear of misunderstanding, we will, in the following, denote the underlying exponent system $(k, k_0, k_1, \dots)$ by (k) for short, and we will denote the corresponding character simply by

$$\Omega^{(k)}(r).$$

Here, again, to each system of values $(k, k_0, \dots)$ and each unit r there clearly corresponds a character $\Omega^{(k)}(r)$.⁵⁶ [66, p. 445]

In contrast, say, to Hadamard, the index (k) is not arbitrary; rather, (k) is taken to range over the specific representing data. But the notation and organization of the proof enables us to ignore the details of the representation where they are not needed. Kronecker noted immediately that $\Omega^{(k)}(r)$ depends only on the value of r modulo m , since any two values with the same residue have the same index. He also notes that we have $\Omega^{(k)}(rr') = \Omega^{(k)}(r)\Omega^{(k)}(r')$. The choice of representation has the further nice property that $\Omega^{(k)}(r)\Omega^{(k')}(r) = \Omega^{(k+k')}(r)$ where $(k+k')$ denotes the result of adding the elements of the tuples (k) and (k') , modulo the cardinality of the associated cyclic groups. Kronecker presented the first orthogonality principle:

$$\sum_{(r)} \Omega^{(0)}(r) = \varphi(m),$$

where r ranges over a system of residues of the units modulo m and (0) is the index of the trivial character, and

$$\sum_{(r)} \Omega^{(k)}(r) = 0$$

for the remaining characters. Kronecker's proof requires unfolding the notation and calculating, but thereafter the fact can be recalled and used in the above form. Similarly, he expressed the second orthogonality relation by the equations

$$\sum_{(k)} \Omega^{(k)}(r_0) = \varphi(m),$$

when r_0 is congruent to 1 modulo m , and

$$\sum_{(k)} \Omega^{(k)}(r) = 0$$

⁵⁶ "Wenn kein Mißverständnis zu befürchten ist, wollen wir im folgenden das zu Grunde gelegte Exponentensystem $(k, k_0, k_1, \dots)$ kurz durch (k) und den zugehörigen Charakter einfacher durch

$$\Omega^{(k)}(r)$$

bezeichnen. Auch hier entspricht für ein festes Wertsystem $(k, k_0, \dots)$ jeder Einheit offenbar r ein Character $\Omega^{(k)}(r)$."

otherwise. He also expressed the dependence of a Dirichlet series on a character in terms of a dependence on (k) , by writing

$$L^{(k)}(z) = \sum_{n=1}^{\infty} \frac{\Omega^{(k)}(n)}{n^z} = \prod_p \frac{1}{1 - \frac{\Omega^{(k)}(p)}{p^z}}.$$

Kronecker’s presentation provides us with an important lesson. The exposition is clearly designed to bring the central ideas to the fore and highlight relevant information, but succeeds in doing so while providing explicit representations and algorithms throughout. This shows that although the moves towards abstraction that we have documented in this section can sometimes *make it possible* to suppress or even ignore algorithmic information, they do not *require* one to do so. In other words, the conceptual reorganization opens the door to the use of other means of describing mathematical objects and operations on them, means that can supplant algorithmic aspects. The question as to whether such means are permissible, meaningful, and appropriate to mathematics lay at the core of twentieth century foundational debates.

7.5 Landau

Born in 1877, Edmund Landau received his doctorate at the University of Berlin in 1899, having studied number theory under Frobenius. He completed a *Habilitation* thesis on Dirichlet series in 1901. Landau later presented proofs of Dirichlet’s theorem in two textbooks, his 1909 *Handbuch der Lehre von der Verteilung der Primzahlen* [68] and his 1927 *Vorlesungen über Zahlentheorie* [69]. We shall here focus on his presentation in the 1909 work, since the later presentation is already essentially what we have portrayed as “contemporary” in Section 4.

Landau began by introducing the characters via the construction in terms of primitive roots and roots of unity. However, the notation that he used to denote them changed over the course of the book. Initially, he used a notation that is quite similar to Dirichlet’s notation for L -functions, writing

$$\chi_{(a_1, a_2, \dots, a_r, a, b)}(n),$$

where the index system $a_1, a_2, \dots, a_r, a, b$ serves to distinguish the characters just as Dirichlet’s $\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{c}', \dots$ distinguished his L -functions. However, after proving that there are $\varphi(d)$ such characters, he simplified his notation to

$$\chi_1(n), \chi_2(n), \dots, \chi_{\varphi(d)}(n),$$

and to $\chi_x(n)$ in the general case, thus adopting notation similar to both de la Vallée-Poussin and Hadamard.

But whereas Landau constructed the characters in the same way as Dirichlet, Dedekind, Hadamard and de la Vallée-Poussin, he also recognized that they are characterized by certain key *properties*, and that it is only these properties that are needed in the proof. Indeed, after introducing the abbreviations for the characters, Landau wrote:

... I will prove four short and elegantly-worded theorems about them [the characters]. The reader may then quickly forget the rather complicated definition of these functions completely, and need only remember that the existence of a system of h distinct functions which possesses the four properties has been proved.⁵⁷ [68, p. 404]

The four theorems that Landau was referring to are the following:

Theorem 1: For any two positive numbers n and n' ,

$$\chi(nn') = \chi(n)\chi(n').$$

This “law of multiplication” holds for each of the h functions ...

Theorem 2: For $n \equiv n' \pmod{k}$,

$$\chi(n) = \chi(n')$$

...

Theorem 3: When n runs through a complete residue system modulo k , for $x = 1$, i.e. for the principal character

$$\sum_n \chi_x(n) = h,$$

however for $x = 2, \dots, h$, i.e. for all other characters

$$\sum_n \chi_x(n) = 0$$

...

Theorem 4: When n is fixed and the sum

$$\sum_{x=1}^h \chi_x(n)$$

extends over all h functions, then

$$\sum_{x=1}^h \chi_x(n) = h \text{ for } n \equiv 1 \pmod{k},$$

$$\sum_{x=1}^h \chi_x(n) = 0 \text{ for } n \not\equiv 1 \pmod{k},$$

therefore for all $k-1$ other residue classes modulo k .⁵⁸ [68, pp. 401–408]

⁵⁷ “...ich werde über sie vier Sätze mit sehr kurzem und elegantem Wortlaut beweisen. Alsdann darf der Leser bald die recht komplizierte Definition dieser Funktionen vollkommen vergessen und braucht sich nur zu merken, daß die Existenz eines Systems von h verschiedenen Funktionen bewiesen worden ist, welche die vier Eigenschaften besitzen.”

⁵⁸ “**Satz 1:** Es ist für zwei ganze positive Zahlen n, n'

$$\chi(nn') = \chi(n)\chi(n').$$

These are exactly the four “important relations” given by de la Vallée-Poussin. What is novel here is Landau’s explicit recognition that only these properties are used in the proof, and that the specific construction is only needed to show the existence of a system of functions that satisfy them.

Like Hadamard, Landau did not go so far as to allow the characters themselves to index the sum in theorem Theorem 4 above, relying on a natural number proxy. But, like de la Vallée-Poussin, he gave an extensional classification of the three types of characters, though he also gave an additional description of the real characters in intensional terms.

Just as Landau’s notation for characters changed over the course of the book, so, too, did his notation for L -functions. Initially, he denoted them in a manner similar to Hadamard’s, writing $L_x(s) = \sum_{n=1}^{\infty} \frac{\chi_x(n)}{n^s}$. However, later on, after concluding his presentation of the proof of Dirichlet’s theorem, Landau adopted a “more convenient” notation:

Now let

$$L_x(s) = \sum_{n=1}^{\infty} \frac{\chi_x(n)}{n^s}$$

be the function corresponding to the character $\chi(n) = \chi_x(n)$; it is now more convenient to include the character in the notation,

$$L(s, \chi),$$

Von jeder der h Funktionen wird also dies “Multiplikationsgesetz” behauptet ...

Satz 2: Es ist für $n \equiv n' \pmod{k}$

$$\chi(n) = \chi(n')$$

...

Satz 3: Wenn n ein vollständiges Restsystem modulo k durchläuft, ist für $x = 1$, d.h. für den Hauptcharakter

$$\sum_n \chi_x(n) = h,$$

dagegen für $x = 2, \dots, h$, d.h. für alle übrigen Charaktere

$$\sum_n \chi_x(n) = 0$$

...

Satz 4: Wenn n festgehalten und die Summe

$$\sum_{x=1}^h \chi_x(n)$$

über alle h Funktionen erstreckt wird, so ist

$$\sum_{x=1}^h \chi_x(n) = h \text{ für } n \equiv 1 \pmod{k},$$

dagegen

$$\sum_{x=1}^h \chi_x(n) = 0 \text{ für } n \not\equiv 1 \pmod{k},$$

also für alle $k - 1$ übrigen Restklassen modulo k .”

and, only when there is no fear of misunderstanding, write

$$L(s)$$

for short, as before.⁵⁹ [68, p. 482]

Landau did not explain *why* the modern notation is more convenient, but we will offer some suggestions in the next section.

Landau’s 1909 presentation is interesting because it has a very transitional feel. Although he gave an intensional construction of the characters, he not only provided a thorough enumeration of their properties, but went out of his way to emphasize that these properties are all that matters to the proof. This is borne out by the fact that the usual division of characters into three classes can be carried out extensionally. And although he initially introduced natural number indices for the characters and L -functions and summed over these indices, he eventually adopted the modern functional notation for L -functions.

When he wrote his 1927 textbook [69], Landau finally made the transition to a proof that is extremely close to the contemporary version. Indeed, he no longer defined the characters by describing how they are constructed, but, rather, defined them in terms of their characteristic properties. Moreover, he summed over the characters themselves, using expressions such as $\sum_{\chi} \chi(a)$. The only real difference between his proof and the one we presented in Section 4.4 is that he did not develop the general notion of a group-theoretic character, but, rather, defined them in terms of the *particular* group of units modulo m . This makes sense, given that the work is an elementary number theory textbook and characters are needed for any other purpose. Roughly speaking, if we combine Landau’s 1927 presentation with Weber’s 1882 more general treatment of characters, the result the presentation in Section 4.

8 Mathematical concerns

In Section 5.2, we considered a number of ways in which characters are treated as bona-fide objects in contemporary proofs of Dirichlet’s theorem. We noted that none of these features are present in Dirichlet’s original proof, for the simple reason that Dirichlet did not isolate or identify the notion of a character.

⁵⁹ “Es sei nun

$$L_x(s) = \sum_{n=1}^{\infty} \frac{\chi_x(n)}{n^s}$$

die dem Charakter $\chi(n) = \chi_x(n)$ entsprechende Funktion; es ist jetzt bequemer, um die Charakter in die Bezeichnung aufzunehmen,

$$L(s, \chi)$$

zu schreiben, und nur, wenn kein Mißverständnis zu befürchten ist, wie früher kurz

$$L(s)."$$

Dedekind's 1863 presentation of Dirichlet's theorem did not use the term "character," but he did introduce the notation, $\chi(n)$, for characters, and identified the defining property of a homomorphism as their "characteristic property." As noted in Section 7.1, after Weber's paper of 1882 the general notion of a group character was in place, and all the subsequently published proofs of Dirichlet's theorem use the terminology of characters.

But now the set of characters modulo m can be defined extensionally, as the set of nonzero homomorphisms from $(\mathbb{Z}/m\mathbb{Z})^*$ to the complex numbers, or intensionally, as functions defined by certain algebraic expressions involving certain primitive elements modulo the prime powers occurring in the factorization of m , and certain complex roots of unity. Even though the two definitions give rise to the same set of characters, proofs can differ in the extent to which they rely on the specific representations or the abstract characterizing property. Dirichlet's proof relied only on the symbolic representations. Somewhat surprisingly, both Dedekind's and Hadamard's division of the characters into the trivial, real, and complex cases was also described in terms of the characters' representations, even though the distinction is naturally expressed in terms of the values they take. Kronecker and de la Vallée-Poussin provided both descriptions, and even though Kronecker made it clear that all operations and classifications can be carried out, algorithmically, in terms of the canonical representations, his careful choice of notation and organization made the extensional properties salient. By 1927, Landau clearly favored the extensional characterization in his textbook.

We have also observed that modern notation like $\sum_{\chi} \chi(n)$ allows us to carry out summations over the finite set of characters modulo m , but that this notation was not available to Dirichlet's early expositors. Dirichlet, Dedekind, and Kronecker all took summations to range over the tuples of integers representing the characters via an explicit algebraic definition, though Kronecker's way of letting a variable (r) range over these tuples is more attractive than Dirichlet's use of $\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{c}', \dots$. In 1882, Weber wrote his sums with ellipses. Curiously, Hadamard in 1896, and Landau in 1909, assigned arbitrary integer indices to the characters, and took sums to range over those indices. The only nineteenth century author to take summation over characters at face value was de la Vallée-Poussin, who nonetheless introduced separate notation S_{χ} to denote such sums. By 1927, however, Landau had adopted the modern notation.

Finally, we have emphasized the modern tendency to represent the dependence of an L -series on a character χ as a functional dependence, with the notation $L(s, \chi)$. Once again, de la Vallée-Poussin was the only nineteenth century expositor of Dirichlet's theorem to do so, with the notation $Z(s, \chi)$. We saw that Landau made the transition in the middle of his 1909 book, and that in his 1927 textbook he relied exclusively on the notation $L(s, \chi)$.

At issue in all these developments is whether characters could be treated in much the same way as natural numbers and real numbers, or whether characters are different sorts of objects, whose treatment has to be mediated by more "concrete" mathematical representations. We contemporary readers of the nineteenth century literature are now so familiar with a modern perspective that it can be hard for us to appreciate the reasons it took so long for the community

to adopt it. Let us begin our analysis of the history, then, by reflecting on the countervailing pressures.

We have had a lot to say about the notational and expressive conventions at play. The difficulty of settling on stable and useful conventions should not be minimized. For example, any mathematician writing a proof has to choose names for variables: should one use m , n , and k to range over natural numbers, or x , y , and z ? One desiderata is to maintain continuity with the background literature, but other constraints come into play: for example, one reason to favor m , n , and k may be that x , y , and z are natural choices to range over other objects, like real or complex numbers, arising in the proof. It is significant, though not surprising, that Dedekind, 26 years after Dirichlet's proof was published, and de la Vallée-Poussin, fully 60 years later, both stuck with Dirichlet's choice of m and k for the first term and common difference in the arithmetic progression, respectively. Even today, q is often used to range over prime numbers in the definition of the L -series, and we still use the letter L exclusively, though most contemporary proofs use a and d in place of m and k .

In a similar way, one has to settle on choices of notation. This also requires thought, even in places where the norms that govern the notation are fairly clear. For example, assuming one divides the set C of characters modulo k into the set consisting of the trivial character, C_{triv} , the set of real characters, C_{real} , and the set of complex characters, $C_{complex}$, it is clear that a sum over C can be broken up accordingly:

$$\sum_{\chi \in C} F(\chi) = \sum_{\chi \in C_{triv}} F(\chi) + \sum_{\chi \in C_{real}} F(\chi) + \sum_{\chi \in C_{complex}} F(\chi).$$

But, even so, one has to settle on the notation to express this relationship, and, as the history of Dirichlet's theorem shows, it can be a long time before a particular means of expression becomes standard. The modern notation emphasizes the parallels between summing over sets C of characters and summing over sets C of natural numbers, but it is by no means obvious that conflating the two is a good idea.

Contemporary readers of Frege's *Begriffsschrift* [43] and *Grundgesetze* [47], which we will discuss in Sections 9 and 10, may be struck by how many pages are devoted to explaining the syntax of the formal language. For those familiar with modern logic, Frege's lengthy explanations seem fiddly and pedantic. But, for Frege, getting the grammatical rules worked out was much of the battle. The syntax of a language only seems trivial when you already know how to speak it.

But it would be a mistake to suggest that all the considerations that Dirichlet's successors faced were "merely" notational. Means of expression often rely on substantial features of our understanding of the nature of that which is expressed. For example, consider the modern way of writing Dirichlet's famous example of 1927:

$$f(x) = \begin{cases} a & \text{if } x \text{ is rational} \\ b & \text{otherwise.} \end{cases}$$

This bears superficial similarity to case-based definitions of number-theoretic functions, or piecewise definitions of analytic functions, that were familiar at the time. But it was novel to base a case distinction on the property of being rational, and, indeed, the notation masks significant assumptions about what it means to define a function on the reals. Consider Frege’s brief account of the evolution of the function concept in 1891:

In the first place, the field of mathematical operations that serve for constructing functions has been extended. Besides addition, multiplication, exponentiation, and their converses, the various means of transition to the limit have been introduced—to be sure, people have not always been clearly aware that they were thus adopting something essentially new. People have gone further still, and have actually been obliged to resort to ordinary language, because the symbolic language of Analysis failed; e.g. when they were speaking of a function whose value is 1 for rational and 0 for irrational arguments.⁶⁰ [45, p. 12]

Frege was not merely concerned to have a convenient notation to express the definition by cases. The definition of f above does that perfectly well, even today. The point is rather that the notation should come with clear rules of use. That is what Frege took to be lacking in the casual use of ordinary language, and what he took his formal system to provide. Similar methodological concerns lie beneath the surface whenever we write $\lim_n a_n$ to denote the limit of a sequence of real numbers, or $I + J$ to denote the sum of two ideals in a ring of algebraic integers. There is nothing tame about the infinitary operations that underlie the notation.

In contrast, foundational questions regarding the use of characters may seem mild. After all, it is easy to represent the characters modulo a positive integer k , and any character is determined by the values it takes on the finitely many residues modulo k . Nonetheless, the use and treatment of characters in proofs of Dirichlet’s theorem bears upon central questions regarding the use and treatment of functions more generally, specifically regarding the relationship between a function and its various representations.

Let us think of Dirichlet L -functions $L(s, \chi)$ in quasi-computational terms. Such a function should take, as input, a real (or complex) value s and a character χ , and return a complex number. Let us set aside the (important) question as to what it means to take a real or complex number as input, or return one as output. What does it mean to accept a character as an input? Should one think of the character as being “presented” to the function as an infinite set of input-output pairs? Or as the list of finite values on the residues modulo k ? (In that

⁶⁰ “Erstens nämlich ist der Kreis der Rechnungsarten erweitert worden, die zur Bildung einer Funktion beitragen. Zu der Addition, Multiplikation, Potenzierung und deren Umkehrungen sind die verschiedenen Arten des Grenzüberganges hinzugekommen, ohne daß man allerdings immer ein klares Bewußtsein von dem wesentlich Neuen hatte, das damit aufgenommen werde. Man ist weiter gegangen und sogar genötigt worden, zu der Wortsprache seine Zuflucht zu nehmen, da die Zeichensprache der Analysis versagte, wenn z.B. von einer Funktion die Rede war, deren Wert für rationale Argumente 1, für irrationale 0 ist.”

case, does the same conception work for function arguments that are not finitely determined?) Should one think, rather, of χ as being some sort of procedure, or subroutine? If so, what sorts of procedures and subroutines are allowed in the definition of a functional with argument χ ? Perhaps, instead, we should identify χ , as Dirichlet did, with its representation in terms of an expression involving certain roots of unity. But recall that those representations relied on choices of primitive elements in the representation of k , although it turns out that the value of the L -function does not vary with different choices of the primitive elements. On our conception, does L somehow “depend” on the choices of primitive elements?

These issues arise not only with respect to functional notation, but also with respect to statements involving quantifiers over characters. For example, the proof of the second orthogonality lemma requires the following fact:

If n is relatively prime to k , there is a character χ such that $\chi(n) \neq 1$.

If we prove this as a separate lemma, we can then invoke the lemma for a given k to “obtain” a χ with the relevant property. But what exactly have we “obtained”? A table of values? A procedure? A representation? If the lemma constructs a χ via a representation of one sort, but our proof of the second orthogonality lemma relies on a different representation of characters, is it legitimate to apply the lemma in that context?

The modern theory of computability and the semantics of programming languages offers various ways of thinking about computer programs which take functions as arguments. The issues are subtle and complex. In contrast, modern mathematics followed a route whereby these subtleties are for the most part set aside. In particular, they are deemed incidental to the proof of Dirichlet’s theorem. Roughly speaking, to make sense of a functional $F(f)$ with function argument f , set theory identifies f “canonically” with its extension, a set of input-output pairs, without concern as to how f is represented or how (and whether) one can “compute” $F(f)$. Set theory then imposes on mathematical language the restriction that the definition of such a functional F can only depend on the extension of f . (Or, put differently: modern mathematical conventions evolved to ensure the latter fact, and axiomatic set theory was designed to model and explain those conventions.)

In the nineteenth century, the answers to the questions raised above were not at all obvious. Indeed, they are still debated among logicians and foundationally-minded mathematicians today. Even for those inclined to dismiss those questions as irrelevant to the proof of Dirichlet’s theorem, it would not have been immediately clear as to whether they really could be dismissed, and, if so, how that should be done. What may seem to be “merely notational” developments in the presentation of Dirichlet’s theorem were part and parcel of the broader mathematical community’s attempt to fashion an understanding of functions as objects that would better support the mathematics of the time.

Setting aside issues of meaning, there may also be concerns about *correctness*. Any choice of notation that draws on an analogy between different domains presupposes that the analogy is appropriate, which is to say, that sufficiently

many properties carry over, and that there are sufficient safeguards to bar the ones that do not. For example, a sum of the form $\sum_{x \in S} t(x)$ generally makes sense when S is finite and the term $t(x)$ takes values on a domain with an addition that is associative and commutative, so that the order the one “runs through” the elements of S does not matter. Summing over infinite totalities is more subtle; it typically depends on having a notion of convergence for the domain in which $t(x)$ takes values, and worrying about the order in which terms are summed. Now, choosing the notation $\sum_{\chi} t(\chi)$ may not be a good idea if it encourages users to transfer facts and properties between domains in an invalid way. This would explain why de la Vallée-Poussin chose a special notation S_{χ} for such sums, as such a new symbol would not come with any unwanted baggage. Adopting the contemporary notation may have required some kind of assurance that the intended contexts would be sufficient to control for proper use.

In sum, mathematical conventions evolve, expand, and change. Any time a mathematician writes a line of text, he or she is situated in a tradition with implicit norms and conventions, and the line of text just written becomes part of that tradition. The fact that so many of these conventions are communicated implicitly does not make them any less important to mathematical understanding.

When a mathematician writes a proof, the intent is that the inferences contained therein will be deemed by his or her colleagues to be correct and justified. Where the inferences are instances of familiar patterns, one desires that they will be easily recognized as such, so that the reader’s effort is conserved for more substantial cognitive tasks. When the inferences rely on assumptions that may be considered dubious, or push familiar patterns of reasoning into unexplored territory, there is greater concern not only as to whether the reasoning will be recognized as *correct*, but also as to whether it will be deemed *appropriate* to addressing the mathematical issues at hand. Thus there are always strong pragmatic pressures to stick close to established convention, and one should not expect fundamental aspects of the language of method of mathematics to change in novel ways, unless there are strong forces pushing for such change.

In the case of Dirichlet’s theorem, we have seen some of the ways in which Dirichlet’s successors chose to modify, or “improve,” his presentation. Now let us try to understand some of the perceived benefits. Kronecker’s proof was explicitly designed to fill in information that was absent from Dirichlet’s proof, in the form of explicit bounds on a quantity asserted to exist. Many of the other developments were explicitly designed towards paving the way to useful generalizations. This was clearly Dedekind’s intent, for example, in pointing out that the Euler product formula depended only on certain multiplicative properties of the terms occurring in the sum. Similarly, abstracting proofs of properties of characters on $(\mathbb{Z}/m\mathbb{Z})^*$ that rely on features specific to the integers modulo m paves the way to extending these properties to group characters more generally. Characters and their properties form the basis for representation theory, which has been an essential part of group theory since the turn of the twentieth century [56, 74]. Authors like Hadamard, de la Vallée-Poussin, and

Landau were also interested in extending Dirichlet's methods to other kinds of Dirichlet series, which now play a core role in analytic number theory.

But it would be a mistake to attribute all the benefits of the expository innovations we have considered to increased generality. After all, these innovations play an equally important role in fostering a better understanding of Dirichlet's proof itself, by highlighting key features of the concepts and objects in question, motivating the steps of the proof, and reducing cognitive burden on the reader by minimizing the amount of information that needs to be kept in mind at each step along the way. It is true that these benefits often support generalization, but they do so in part by making our thinking *vis-à-vis* Dirichlet's proof itself more efficient.

For example, proofs of Dirichlet's theorem that rely on explicit representations of the characters require us to keep the details of the representation in mind. Recall Dirichlet's original representation of an arbitrary character:

$$\chi(n) = \theta^\alpha \varphi^\beta \omega^\gamma \omega'^{\gamma'} \dots$$

Here the reader has to keep in mind that $\alpha, \beta, \gamma, \gamma'$, and so on are the indices of n with respect to primitive elements chosen in the decomposition of the abelian group $(\mathbb{Z}/m\mathbb{Z})^*$, and $\theta, \varphi, \omega, \omega'$, and so on are corresponding roots of unity. This information has to be kept in mind throughout the proof, because the nature of the objects and the dependences of $\alpha, \beta, \gamma, \gamma'$ on n may play a role in licensing an inference or calculation. Moreover, recall that we obtain all the characters by expressing all the roots

$$\theta = \Theta^{\mathfrak{a}}, \varphi = \Phi^{\mathfrak{b}}, \omega = \Omega^{\mathfrak{c}}, \omega' = \Omega'^{\mathfrak{c}'}, \dots,$$

in terms of primitive roots of unity $\Theta, \Phi, \Omega, \Omega', \dots$, and letting $\mathfrak{a}, \mathfrak{b}, \dots$ range over the appropriate exponents. Once again, this information has to be remembered throughout. Later proofs are easier to read simply because they do not require us to keep as much information in mind, and highlight the relevant dependences when they are needed.

One way of achieving this is by reorganizing the proof in such a way that some of the relevant information is localized to particular facts and calculations. For example, even if one resorts to representations to prove the orthogonality relations, if this is the only place they are used, then they do not need to be ready to hand when these relations are invoked in a calculation later on, where other analytic expressions and their properties are the objects of focus. Thus modularity reduces cognitive burden, and makes it easier to keep track of the global structure of the argument, providing high-level outlines, or sketches, of the proof. Such restructuring paves the way to generality: isolating context-specific details in well-insulated modules means that one can adapt the proof by changing the modules while preserving their external interfaces. But, to repeat, generality is not the only benefit: the restructuring improves the readability of the original proof as well.

Even in situations where there is a lot of information in play at once, judicious notation and means of expression can make important relationships between the data more salient. For example, in Section 6, we saw that Dirichlet

wrote

$$\begin{aligned} \sum \frac{1}{q^{1+\rho}} + \frac{1}{2} \sum \frac{1}{q^{2+2\rho}} + \frac{1}{3} \sum \frac{1}{q^{3+3\rho}} + \dots \\ = \frac{1}{K} \sum \Theta^{-\alpha_m \mathfrak{a}} \Phi^{-\beta_m \mathfrak{b}} \Omega^{-\gamma_m \mathfrak{c}} \Omega^{-\gamma'_m \mathfrak{c}'} \dots \log L_{\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{c}', \dots} \end{aligned}$$

where we would write

$$\sum_{\chi \in (\widehat{\mathbb{Z}/q\mathbb{Z}})^*} \overline{\chi(m)} \log L(s, \chi) = \varphi(q) \sum_{\substack{p \equiv m \\ (\text{mod } q)}} \frac{1}{q^s} + O(1)$$

The second presentation highlights the relationship between the series $L(s, \chi)$ and the characters; in particular, the value $L(s, \chi)$ is multiplied by the value of the conjugate character, $\overline{\chi}$, at m in each term of the sum on the left-hand side. Dirichlet's notation obscures this relationship, since the logarithm of the appropriate L -series $L_{\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{c}', \dots}$ is multiplied by the character $\Theta^{-\alpha_m \mathfrak{a}} \Phi^{-\beta_m \mathfrak{b}} \Omega^{-\gamma_m \mathfrak{c}} \Omega^{-\gamma'_m \mathfrak{c}'} \dots$ and the reader must sift through this expression to notice that the exponents $-\mathfrak{a}, -\mathfrak{b}, -\mathfrak{c}, -\mathfrak{c}', \dots$ correspond to the subscripts of the L -series.

The same problem applies to Hadamard's method of using arbitrary indices for the characters: given a character ψ_v , there is no natural name for the index of its conjugate $\overline{\psi}_v$. Of course, one can introduce such a notation, but that requires keeping track of the relationship between the two notations, that is, the conjugation of characters and the associated operation on indices. And grouping the characters into different classes requires grouping the indices into different classes, again yielding an uncomfortable duality. (At least Kronecker maintained a monist consistency, insisting that all operations on characters are operations on their representations. So when $\Omega^{(k)}$ is a character, Kronecker could write $\Omega^{(-k)}$ for its conjugate, since the conjugate character is obtained by negating the elements of the corresponding tuple.)

Similar considerations may explain why Landau changed his notation for L -functions in the middle of his 1909 work, from $L_x(s)$ to $L(s, \chi)$. In the paragraph following his notation change, Landau showed that the theory of L -functions can be reduced to L -functions that correspond to particular types of characters, called *proper characters*. Roughly, proper characters modulo k are those which cannot be obtained as a character modulo K where $K < k$. To show that the theory of L -series can be reduced in the appropriate way, he proved that for an improper character modulo k and $\sigma > 1$, we have [68, 482-483]:

$$L(s, \chi) = \prod_{\nu=1}^c \left(1 - \frac{\varepsilon_\nu}{p_\nu^s} L_0(s, X) \right).$$

Here $L_0(s, X)$ is an L -series corresponding to the proper character X modulo K , ε_ν are certain roots of unity, and c a number (which can be 0, depending on how many prime factors of k are contained in K). If Landau had kept his original notation, the left hand side of the above equation would be written as $L_x(s)$

where $\chi = \chi_x$. But how would we represent the right hand side? We should note that one obvious choice for an index, x' , would not be available, since Landau used this previously in connection with the conjugates of characters. In the text, Landau had identified a relationship between a character, χ , and the corresponding proper character, X ; having to translate this to a relationship between indices would only clutter the exposition. These issues are compounded when, in later sections, Landau wanted to obtain functional equations to relate $L(s, \chi)$ and $L(1-s, \overline{\chi})$ and in doing so referred to the distinction between proper and improper characters (see e.g. [68, §130]). Using the old notation, we would need to keep track of three different subscripts to index the various characters.

Consider, finally, the issue of uniformity of notation. We have already discussed concerns associated with the notation $\sum_{\chi}$ for summation over characters. But there is an obvious benefit to using the same notation for summing over finite sets of characters and summing over finite sets of integers, namely, that the two operations really do share common properties. Indeed, the transfer is immediate via the Hadamard trick of assigning an integer index to each character. The option of introducing a new symbol every time one needs to index sums by finite sets of objects is clearly untenable, as it would result in a confusing explosion of notations. It does not seem at all surprising that de la Vallée's notation S_{χ} was short-lived.

To sum up, then, we have identified a number of advantages to the rewritings of Dirichlet's proof considered in the last section:

- Essential properties of key objects (or expressions) were isolated, reducing the amount of information that someone reading the proof has to keep in mind at each step.
- The organization of proofs became more modular, with information localized to very specific parts of the proof.
- Expressions became more readable, since irrelevant details were suppressed, and the features that remained made dependences and relationships between terms more salient.
- Notation became more uniform, highlighting commonalities between different domains.

Changes like this often go hand in hand with attempts to generalize concepts and methods to other domains, since managing and controlling the volume of domain-specific detail tends to bring to the fore aspects of the proof that transcend these specifics. But they also contribute to a better understanding of Dirichlet's proof itself, and make the proof easier to read and reproduce from memory.

Benefits such as these are often dismissed as merely “pragmatic” or “cognitive,” but this downplays the fact that such considerations effectively shape and justify the norms that guide our mathematical practice. The philosophy of mathematics needs to take them seriously.

9 Function and object in Frege

In 1940, Alonzo Church presented a formulation of type theory [17], now known as “simple type theory.” Simple type theory can serve as a foundation for a significant portion of mathematics, and, indeed, is the axiomatic foundation of choice for a number of computational interactive theorem provers today [51, 54, 82]. One starts with some basic types, say, a type $\mathbb{B}$ of Boolean truth values and a type $\mathbb{N}$ of natural numbers, and one forms more complex types $\sigma \times \tau$ and $\sigma \rightarrow \tau$ from any two types σ and τ . Intuitively, objects of type $\sigma \times \tau$ are ordered pairs, consisting of an object of type σ and an object of type τ , and objects of type $\sigma \rightarrow \tau$ are functions from σ to τ . In a type-theoretic approach to the foundations of mathematics, one identifies sets of natural numbers with predicates, which is to say, objects of type $\mathbb{N} \rightarrow \mathbb{B}$. Binary relations on the natural numbers are then objects of type $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{B}$, and sequences of natural numbers are objects of type $\mathbb{N} \rightarrow \mathbb{N}$. Objects at this level are called *type 1* objects, because they require one essential use of the function space arrow. Integers can be identified as pairs of natural numbers and rationals can be identified as integers in the usual ways. Real numbers are then Cauchy sequences of rationals (type 1 objects), or equivalence classes of such, which puts them at type 2. Functions from the reals to the reals and sets of reals are then objects of type 3, and sets of functions from the reals to reals or collections of sets of real numbers are then objects of type 4. For example, the collection of Borel sets of real numbers is a type 4 object, as is Lebesgue measure, which maps certain sets of real numbers to the real numbers. A set of measures on the Borel sets of the real numbers is a type 5 object. And so on up the hierarchy.

Simple type theory can be viewed as a descendent of the ramified type theory of Russell and Whitehead’s *Principia mathematica* [92], which, in turn, was inspired by the formal system of Frege’s *Grungesetze der Arithmetik* [47]. Starting with a basic type of individuals, Frege’s system also has variables ranging over higher-type functionals, and so can be seen as an incipient form of modern type theory. For that reason, it can then come as a surprise to logicians familiar with the modern type-theoretic understanding that the foundational outlook just described is *not at all* the image of mathematics that Frege had in mind. It is this image that we wish to explore here.

Frege took concepts to be instances of functions; for example, in “Function and concept” he wrote that “a concept is a function whose value is always a truth value” [45, p. 139].⁶¹ And, throughout his career, he was insistent that functions are not objects. The third “fundamental principle” in his *Grundlagen der Arithmetik* of 1884 was “never to lose sight of the distinction between concept and object”⁶² [44, Introduction], and he later asserted that “it will not do to call a general concept word the name of a thing” [44, § 51].⁶³ The distinc-

⁶¹We should note that in this section we will focus on his views from 1884 onwards. Prior to this, he seems to have held a different view of concepts, though he still maintained that they are not objects; see [58, p. 136].

⁶²“... der Unterschied zwischen Begriff und Gegenstand ist in Auge zu behalten.”

⁶³“... ist es unpassend, ein allgemeines Begriffswort Namen eines Dinges zu nennen.”

tion features prominently in his essays “Function and Concept,” “Comments on *Sinn* and *Bedeutung*” and “Concept and object” of 1891, 1891/2, and 1892, respectively.

Frege’s distinction was primarily linguistic: objects are denoted by words and phrases that can fill the subject role in a grammatical sentence, whereas concepts are denoted by words and phrases that can play the role of a predicate. In “Concept and object” he wrote:

We may say in brief, taking “subject” and “predicate” in the linguistic sense: a concept is the *Bedeutung* of a predicate; an object is something that can never be the whole *Bedeutung* of a predicate, but can be the *Bedeutung* of a subject.⁶⁴ [46, pp. 198]

And:

A concept—as I understand the word—is predicative. On the other hand, a name of an object, a proper name, is quite incapable of being used as a grammatical predicate.⁶⁵ [46, pp. 193]

In the sentence, “Frege is a philosopher,” the word “Frege” denotes an object, and the phrase “is a philosopher” denotes a concept. Frege clarified the distinction by explaining that functional expressions, including concept expressions, are “unsaturated,” or incomplete. These stand in contrast to signs that are used to denote objects, which are complete in and of themselves. For example, in the sentence “Frege is a philosopher,” the expression “Frege” is saturated, and succeeds in picking out an object. In contrast, the expression “...is a philosopher” contains a gap, and fails to name an object until one fills in the ellipsis, at which point the expression denotes a truth value.⁶⁶

Having distinguished between concepts and objects in such a way, Frege had to deal with objections, such as the one he attributed to Benno Kerry in “Concept and object.” In the sentence “The concept ‘horse’ is a concept easily attained” the concept denoted by ‘horse’ does fill the subject role. Frege’s surprising answer was to deny that the phrase “the concept ‘horse’ ” denotes a concept. He conceded that this sounds strange:

⁶⁴ “Wir können kurz sagen, indem wir “Prädikat” und “Subjekt” im sprachlichen Sinne verstehen: Begriff ist Bedeutung eines Prädikates, Gegenstand ist, was nie die ganze Bedeutung Prädikates, wohl aber Bedeutung eines Subjekts sein kann.” The word *Bedeutung* is often translated as “reference” or “denotation.” But for difficulties in the translation, see §, 4 of the introduction to Beaney [8].

⁶⁵ “Der Begriff—wie ich das Wort verstehe—ist prädikativ. Ein Gegenstandsname hingegen, ein Eigenname ist durchaus unfähig, als grammatisches Prädikat gebraucht zu werden.”

⁶⁶ While the distinction between saturated and unsaturated expressions is cast as a distinction between linguistic signs, in his 1904 essay “What is a Function?” Frege made it clear that the dichotomy extends to functions and objects themselves: “The peculiarity of functional signs, which we here called ‘unsaturatedness’, naturally has something answering to it in the functions themselves. They too may be called ‘unsaturated’ ...” (“Der Eigentümlichkeit der Funktionszeichen, die wir Ungesättigkeit genannt haben, entspricht natürlich etwas an den Funktionen selbst. Auch diese können wir ungesättigt nennen ...”) [48, p. 665].

It must indeed be recognized that we are confronted by an awkwardness of language...if we say that the concept *horse* is not a concept...⁶⁷ [46, pp. 196–197]

Yet, he insisted, this is what we must do. He was already clear about this in the *Grundlagen*:

The business of a general concept word is precisely to signify a concept. Only when conjoined with the definite article or a demonstrative pronoun can it be counted as the proper name of a thing, but in that case it ceases to count as a concept word. The name of a thing is a proper name.⁶⁸ [44, §51]

And so, in “Concept and object,” he reminded us:

If we keep it in mind that in my way of speaking expressions like “the concept *F*” designate not concepts but objects, most of Kerry’s objections already collapse.⁶⁹ [46, pp. 198–199]

He similarly urged us to reconstrue expressions like “all mammals have red blood” as “whatever is a mammal has red blood” so as to avoid the impression that the predicate “has red blood” is being applied to an object, “mammal.” Although these examples deal with concepts, Frege’s analysis makes it clear that he intended the linguistic separation to remain decisive for other kinds of functions as well.

At the same time, Frege was equally dogmatic in insisting that what we commonly take to be mathematical objects really *are* mathematical objects as such. The introduction to his *Grundlagen* begins as follows:

When we ask someone what the number one is, or what the symbol 1 means, we get as a rule the answer “Why, a thing.”⁷⁰ [44, Introduction]

The claim is so curious as to give one pause.⁷¹ The fact that Frege used such a brazen rhetorical flourish to frame the whole project makes it clear just how central the issue is to his analysis. Once again, his reason for maintaining the distinction is largely linguistic; for example, because the number 7 plays the role of a subject in the statement “7 is odd,” 7 must be an object. But, once

⁶⁷ “Es kann ja nicht verkannt werden, daß hier eine freilich unvermeidbare sprachliche Härte vorliegt, wenn wir behaupten: der Begriff Pferd ist kein Begriff...”

⁶⁸ “Ein allgemeines Begriffswort bezeichnet eben einen Begriff. Nur mit dem bestimmten Artikel oder einem Demonstrativpronomen gilt es als Eigenname eines Dinges, hört aber damit auf, als Begriffswort zu gelten. Der Name eines Dinges ist ein Eigenname.”

⁶⁹ “Wenn wir festhalten, daß in meiner Redeweise Ausdrücke wie “der Begriff *F*” nicht Begriffe, sondern Gegenstände bezeichnen, so werden die Einwendungen *Kerry*s schon größtenteils hinfällig.”

⁷⁰ “Auf die Frage, was die Zahl Eins sei, oder was das Zeichen 1 bedeute, wird man meistens die Antwort erhalten: nun, ein Ding.” All our translations from the *Grundlagen* are taken from the Austin translation cited in the references.

⁷¹ We are grateful to Steve Awodey for this observation.

again, Frege had to deal with sentences where the syntactic role of a number is murkier. For example, he considered uses of number terms in language that are attributive and do not occur prefixed by the definite article, for example, “Jupiter has four moons” [44, §57]. He wrote

“... our concern here is to arrive at a concept of number usable for the purposes of science; we should not, therefore, be deterred by the fact that in the language of everyday life number appears also in attributive constructions. That can always be got round.”⁷² [44, §57]

Specifically, it can be got round by writing an attributive statement such as “Jupiter has four moons” as “the number of Jupiter’s moons is the number 4, or 4” [44, §57], thereby eliminating the attributive usage.

So, for Frege, functions are not objects, but numbers are, because they play the subject role in mathematical statements and can be used with the definite article. There is clearly a difficulty lurking nearby. At least from a modern standpoint, we tend to view functions, sequences, sets, and structures as objects, and certainly in Frege’s time locutions such as “the function f ” and “the series s ” were common. Frege’s response was similar to his response to Kerry’s objection, namely, to deny that these expressions like these denote functions. To understand how this works, consider the fact that Frege’s logical system includes an operator which takes any function f from objects to objects and returns an object, $\dot{e}f(\varepsilon)$, intended to denote its “course-of-values” or “value range.” If f is a concept, which is to say, a function which for each object returns values, the course-of-values of f is called the “extension” of the concept. Frege’s Basic Law V asserts that two functions which are extensionally equal—that is, which return equal output values for every input—have the same courses-of-values.

Frege used these courses-of-values and extensions as object-proxies for functions and concepts. This is how he analyzed the concept of a cardinal number. Let F , for example, be a second-level concept, such that F holds of a first-level concept f if and only if f holds of exactly one object. Frege took the number one to be the extension of F , thereby achieving the goal of making the number one, well, a thing. But this “pushing down” trick is central to the methodology of the *Grundgesetze*: whenever the formal analysis of common mathematical objects seems to suggest identifying such objects as functions or concepts, Frege avoided doing so by replacing the function or concept with its extension. For example, in the *Grundgesetze* he circumvented the need to define mathematical operations on sequences and relations construed as functions, defining the operations rather on the associated courses-of-values.⁷³ Referring to Frege’s concepts as

⁷² “Da es uns hier darauf ankommt, den Zahlbegriff so zu fassen, wie er für die Wissenschaft brauchbar ist, so darf es uns nicht stören, dass im Sprachgebrauche des Lebens die Zahl auch attributiv erscheint. Das lässt sich immer vermeiden.”

⁷³ In fact, the definition of the number one earlier in the paragraph describes, more precisely, the construction in the *Grundlagen*. In the *Grundgesetze*, he took F to be a *first-level* concept that holds of classes, i.e. extensions of concepts, that contain one element. This is a nice illustration of how the “pushing down” trick can be used repeatedly to avoid the use of higher types. See Reck [90, Section 5] for a discussion of the two definitions, and Burgess [11] for an overview of Frege’s methodology.

“attributes” and their extensions as “classes,” Quine described the difference as follows:

Frege treated of attributes of classes without looking upon such discourse as somehow reducible to a more fundamental form treating of attributes of attributes. Thus, whereas he spoke of attributes of attributes as *second-level* attributes, he rated the attributes of classes as of first level; for he took all classes as rock-bottom objects on par with individuals. [86, p. 147]

Frege never got so far as developing mathematical analysis in his system, and we cannot say how with certainty how he would have developed, for example, ordinary calculus on the real numbers. But there is a strong hint that here, too, he would have taken, for example, operations like integration and differentiation to operate on extensions, rather than functions, in his system. He touched on the history of analysis in his *Function and concept* of 1891, and noted that, for example, differentiation can be understood as a higher-type functionals.

Now at this point people had particular second-level functions, but lacked the conception of what we have called second-level functions. By forming that, we make the next step forwards. One might think that this would go on. But probably this last step is not so rich in consequences as the earlier ones; for instead of second-level functions one can deal, in further advances, with first-level functions—as shall be shown elsewhere.⁷⁴ [45, p. 31]

Presumably, he had the method of replacing functions by their extensions in mind.

Notice, incidentally, that Frege’s method of representing mathematical functions as courses-of-values has the effect that mathematical functions are treated extensionally. For example, defining the integral as an operation that applies to a course-of-values means that integration cannot distinguish mathematical functions that are extensionally equal, since any two descriptions of a function that satisfy extensional equality have the same course-of-values, by Basic Law V.

There are other interesting features of Frege’s treatment of functions and higher-type objects that push us away from identifying them with the functions of ordinary mathematics. For example, for Frege, every function has to be defined on the entire domain of individuals; even if one is interested in the exponential function on the real numbers, one has to specify a particular (but arbitrary) value of this function for every object in existence. And the separation of functions and objects has other effects on the system. There is only one basic

⁷⁴ “Damit hatte man nun einzelne Funktionen zweiter Stufe, ohne jedoch das zu erfassen, was wir Funktion zweiter Stufe genannte haben. Indem man dies tut, macht man den nächsten Fortschritt. Man könnte denken, dass dies so weiter ginge. Wahrscheinlich ist aber schon dieser letzte Schritt nicht so folgenreich wie die früheren, weil man statt der Funktionen zweiter Stufe im weiteren Fortgang Funktionen erster Stufe betrachten kann, wie an einem anderen Orte gezeigt werden soll.”

type, so, for example, truth values live alongside everything else. There is no notion of identity between higher-type objects—the equality symbol can only be applied to equality between objects—even though Frege pointed out that one can define the notion of “sameness” of functions and concepts extensionally, and noted that this acts very much like identity. Frege’s system, of course, includes the axiom of universal instantiation. In contemporary notation, this would be expressed as $\forall x \varphi(x) \rightarrow \varphi(a)$ where x is a variable ranging over individuals and a is any individual term. It also includes the corresponding axiom $\forall F \varphi(F) \rightarrow \varphi(A)$, where F ranges over functions from objects to objects. But, remarkably, the system does not include analogous axioms for higher-type objects: the “pushing down trick” obviates the need for these.

All things considered, Frege’s foundational treatment of mathematics seems much closer to modern set-theoretic treatments, where there is one homogeneous universe of individuals. Truth values are individuals, numbers are individuals, mathematical sequences and series are individuals—all bona-fide mathematical objects are individuals. Functions play a mere linguistic role in the syntax of the system, providing the axioms and rules enough flexibility to get the job done. They are not the cogs in the system, but, rather, the oil the keeps the wheels turning. As Panza puts it:

... according to Frege, appealing to functions is indispensable in order to fix the way his formal language is to run, but functions are not as such actual components of the language. More generally, functions manifest themselves in our referring to objects—either concrete or abstract—and making statements about them, but they are not as such actual inhabitants of some world of *concreta* and *abstracta*. Briefly: Frege’s formal language, as well as our ordinary ones, display functions, but there are no functions as such. [83, p. 14]

10 Foundational concerns

We have seen that a curious tension lies at the core of Frege’s formal representation of mathematics. On the one hand, Frege asserted, repeatedly, that functions, in the logical and linguistic sense, are not objects. On the other hand, when it comes to formalizing mathematical constructions, he clearly felt that functions, in the mathematical sense, *have to be* objects. His course-of-values operator, together with his Basic Law V, allowed him to have his cake and eat it too, maintaining clear borders between the two realms while passing between them freely. But Frege is often taken to task for failing to realize that this strategy opens the door to Russell’s paradox. Indeed, the strategy feels like a hack, a desperate attempt to satisfy the two central constraints. Why was he so committed to them? The goal of this section is to suggest that the concerns Frege was trying to address with the design of the logic of the *Grundgesetze* parallel some of the informal mathematical concerns we were able to discern in the nineteenth century treatment of characters.

When one considers the philosophical and logical considerations that went into the design of Frege's logic, two possibilities come to mind. One is that Frege determined that functions and objects should be separate on broad metaphysical grounds, and then designed the logic accordingly. The other is that he designed the logic, determined it worked out best with a separation of individuals and functions, and read off the metaphysical stance from that. But, in fact, there is no clear distinction between these two descriptions. Frege designed his logic to try to model scientific practice at its best, and account for and support its successes while combating and eliminating confusions. The examples in the previous section show that Frege had no qualms about reinterpreting ordinary locutions and reconstruing everyday language, so he was by no means slave to naive metaphysical intuitions. But even when doing so he appealed to intuitions to convince us that the reconstructions are reasonable. Thus "doing metaphysics" meant analyzing the practice, sorting out intuitions, and trying to regiment and codify them in a coherent and effective way. From the other direction, "getting the logic to work" meant being able to account for the informal practice effectively and efficiently, and supporting our intuitions to the extent that they can be fashioned into a coherent system. So it is not a question as to whether the metaphysics or the logic comes first; working out the metaphysics and designing the logic are part and parcel of the same enterprise. The following questions therefore seem more appropriate:

1. What considerations pushed Frege to maintain the sharp distinction between function and object?
2. What considerations pushed Frege to identify mathematical entities, including ordinary mathematical functions, as objects?

Let us consider each in turn.

It seems to us that the answer to the first question is simply that Frege felt that failure to respect the distinction results in linguistic confusion.

If it were correct to take "one man" in the same way as "wise man," we should be able to use "one" also as a grammatical predicate, and to be able to say "Solon was one" just as much as "Solon was wise." It is true that "Solon was one" can actually occur, but not in a way to make it intelligible on its own in isolation. It may, for example, mean "Solon was a wise man," if "wise man" can be supplied from the context. In isolation, however, it seems that "one" cannot be a predicate. This is even clearer if we take the plural. Whereas we can combine "Solon was wise" and "Thales was wise" into "Solon and Thales were wise," we cannot say "Solon and Thales were one." But it is hard to see why this should be impossible, if "one" were a property both of Solon and of Thales in the same way that "wise" is.⁷⁵ [44, §29]

⁷⁵ "Wenn 'Ein Mensch' ähnlich wie 'weiser Mensch' aufzufassen wäre, so sollte man denken, dass, 'Ein' auch als Praedicat gebraucht werden könnte, sodass man wie 'Solon war weise'

In other words, even though in some contexts an object word like “one” can *appear* to be used as a predicate, and in other contexts a concept can *appear* to be used as a subject, closer inspection shows that these uses do not conform to the rules that govern the use of prototypical subjects and predicates, and so should not be categorized in the naive way.

One of Frege’s favorite pastimes was to show that assertions made by philosophical and mathematical colleagues degenerate into utter nonsense when they fail to maintain sufficient linguistic hygiene. For example, in his 1904 essay, “What is a Function,” Frege was critical of conventional mathematical accounts of variables and functions. It is a mistake, he said, to think of a variable as being an object that varies:

... a number does *not* vary; for we have nothing of which we could predicate the variation. A cube never turns into a prime number; an irrational number never becomes rational.⁷⁶ [48, p. 658]

He took the mathematician Emanuel Czuber to task for giving such a sloppy account of variables and functions in an introductory mathematical text. For example, he criticized Czuber’s terminology “a variable assumes a number” [48, 288] as being incomprehensible. On Czuber’s account, a variable is an “indefinite number,” so the terminology can be rephrased “an indefinite number assumes a (definite) number”; but where we may talk about an object assuming a property, what can it mean for an object to assume another object?

In other connections, indeed, we say that an object assumes a property, here the number must play both parts; as an object it is called a variable or a variable magnitude, and as a property it is called a value. That is why people prefer the word “magnitude” to the word “number”; they have to deceive themselves about the fact that the variable magnitude and the value it is said to assume are essentially the same thing, that in this case we have *not* got an object assuming different properties in succession, and that therefore there can be no question of a variation.⁷⁷ [48, p. 660–661]

auch sagen könnte ‘Solon war Ein’ oder ‘Solon war Einer’. Wenn nun der letzte Ausdruck auch vorkommen kann, so ist er doch für sich allein nicht verständlich. Er kann z.B. heissen: Solon war ein Weiser, wenn ‘Weiser’ aus dem Zusammenhange zu ergänzen ist. Aber allein scheint ‘Ein’ nicht Praedicat sein zu können. Noch deutlicher zeigt sich dies beim Plural. Während man ‘Solon war weise’ und ‘Thales war weise’ zusammenziehen kann in ‘Solon und Thales waren weise,’ kann man nicht sagen ‘Solon und Thales waren Ein’. Hiervon wäre die Unmöglichkeit nicht einzusehen, wenn ‘Ein’ sowie ‘weise’ eine Eigenschaft sowohl des Solon als auch des Thales wäre”.

⁷⁶ “Folglich verändert sich die Zahl gar nicht; denn wir haben nichts, von dem wir die Veränderung aussagen könnten. Eine Kubikzahl wird nie zu einer Primzahl, und eine Irrationalzahl wird nie rational.”

⁷⁷ “Sonst sagt man wohl, daß ein Gegenstand eine Eigenschaft annehme; hier muß die Zahl beide Rollen spielen; als Gegenstand wird sie Variable oder veränderliche Größe, als Eigenschaft wird sie Wert genannt. Darum also zieht man das Wort ‘Größe’ dem Worte ‘Zahl’ert, den sie angeblich annimmt, im Grunde dasselbe sind, daß man gar nicht den Fall hat, wo ein Gegenstand nacheinander verschiedene Eigenschaften annimmt, daß also von Veränderung in keiner Weise die Rede sein kann.”

The essay closes with the following assessment:

The endeavor to be brief has introduced many inexact expressions into mathematical language, and these have reacted by obscuring thought and producing faulty definitions. Mathematics ought properly to be a model of logical clarity. In actual fact there are perhaps no scientific works where you will find more wrong expressions, and consequently wrong thoughts, than in mathematical ones. Logical correctness should never be sacrificed to brevity of expression. It is therefore highly important to devise a mathematical language that combines the most rigorous accuracy with the greatest possible brevity. To this end a symbolic language would be best adapted, by means of which we could directly express thoughts in written or printed symbols without the intervention of spoken language.⁷⁸ [48, p. 665]

Frege aimed to give a clear account of the rules that govern proper logical reasoning. Although, in ordinary language, the line between concepts and objects is sometimes blurry, failure to diagnose and manage the blurriness open the door to nonsensical reasoning. Even though words like “one” and “horse” sometimes seem to denote both concepts and objects, conflating the two causes problems. For Frege, the only viable solution was to analyze and regiment such uses in a way that cordons off problematic instances. He found that the best way to do this is to maintain a clear separation of concept and object, and then supplement the analysis with an explanation as to how some words seem to cross the divide in certain contexts.

Now let us turn to the second question: why was Frege so dogged in his insistence that mathematical entities like numbers have to be treated as objects, and so persistent, in practice, in pushing mathematical constructions down to that realm? We believe that the answer lies in an observation that we found in Heck [57]: Frege wanted his numbers to be able to count all sorts of entities, and the only way he could make that work was by treating all these entities as inhabitants of the same type. Consider the following statements:

- There are two truth values.
- There are two natural numbers strictly between 5 and 8.
- There are two constant functions taking values among the truth values.

⁷⁸“Das Streben nach Kürze hat viele ungenaue Ausdrücke in die mathematische Sprache eingeführt, und diese haben rückwirkend die Gedanken getrübt und fehlerhafte Definitionen zuwege gebracht. Die Mathematik sollte eigentlich ein Muster von logischer Klarheit sein. In Wirklichkeit wird man vielleicht in den Schriften keiner Wissenschaft mehr schiefe Ausdrücke und infolgedessen mehr schiefe Gedanken finden als in den mathematischen. Niemals sollte man die logische Richtigkeit der Kürze des Ausdrucks opfern. Deshalb ist es von großer Wichtigkeit, eine mathematische Sprache zu schaffen, die mit strengster Genauigkeit möglichste Kürze verbindet. Dazu wird wohl am besten eine Begriffsschrift geeignet sein, ein Ganzes von Regeln, nach denen man durch geschriebene oder gedruckte Zeichen ohne Vermittlung des Lautes unmittelbar Gedanken auszudrücken vermag.”

- There are two characters on $(\mathbb{Z}/4\mathbb{Z})^*$.
- There are two subsets of a singleton set.

Frege would have insisted that the word “two” in each of these statements refers to the same object. We would like to say that the number of truth values is equal to the number of natural numbers between 5 and 8, but if truth values and numbers were different types of entities, his analysis of number would not do that. The problem is that Frege’s grammar requires a different notion of equinumerosity for each pair of types. In other words, for any two types σ and τ , there is an expression $\text{Eq}_{\sigma,\tau}$ that expresses equinumerosity between concepts of entities of those types. Similarly, for each type σ , there is a concept Two_σ that holds of concepts of arguments of type σ under which two elements fall.⁷⁹ Taking extensions of each of these, following Frege’s construction, would yield an object 2_σ for each type σ . But this results in a proliferation of twos, and since 2_σ and 2_τ are not guaranteed to be the same object, one would have to exercise great care when reasoning about the relationship between them. This is clearly unworkable.

Instead, Frege designed his numbers to count objects, *simpliciter*: 2 is the extension of the concept of being a concept of *objects* under which exactly two objects fall.⁸⁰ But this means that if you want to count a collection of things, those things have to be elements of the type of objects. This, in turn, provides a strong motivation to locate mathematical entities of all kinds among the type of objects.

We have seen in Sections 4 to 8 that what holds true of counting holds true of other mathematical operations, relations, and constructions as well. Contemporary proofs of Dirichlet’s theorem have us sum over finite sets of characters just as we sum over finite sets of numbers. We view the general operation here as summation over a finite set of objects, viewing both characters and numbers as such. Contemporary proofs also have us consider groups of characters, just as we consider groups of residues. Once again, we consider these as instances of the general group concept, with the understanding that a group’s underlying set can be any set of objects. This allows us to speak over a homomorphism between any two groups, without requiring a different notion of “homomorphism” depending on the type of objects of the groups’ carriers.

Characters were not the only mathematical entities studied in the latter half of the nineteenth century that encouraged set-theoretic reification. Gauss’ genera of quadratic forms, discussed in Section 7.1, also bear a group-theoretic structure, and these are sets of quadratic forms. Dedekind developed his theory of ideals in order to supplement rings of algebraic integers with “ideal divisors,” extending the unique factorization property of the ordinary integers to these more general domains. Dedekind found that these ideal divisors could be identified with sets of elements in the original ring, now known as “ideals.” Like the

⁷⁹In Frege’s system, for σ anything other than the type of objects, this has to be expressed in terms of a “sameness” relation for elements of type σ , since the equality relation only applies to objects.

⁸⁰See footnote 73.

characters, the ideals of a ring of algebraic integers bear an algebraic structure, and Dedekind was adamant that they should be treated as bona-fide objects.⁸¹ Similarly, Dedekind constructed the real numbers by indentifying each of them with a pair of sets of rational numbers [21]. By the end of the century, it was common to view a quotient group as a group whose elements are equivalence classes, or cosets.⁸²

The reasons given above to treat mathematical functions and sets as objects also speak in favor of treating them extensionally. The statements that “there are two characters on $\mathbb{Z}/2\mathbb{Z}$ ” and “there are $\varphi(m)$ characters on $\mathbb{Z}/m\mathbb{Z}$ ” are false if we take characters to be representations, as there are many different representations of the same character. We could, of course, develop notions of “counting up to equivalence.” In the early days of finite group theory, Camille Jordan described quotient groups as systems just like ordinary groups except that equality is replaced by an appropriate equivalence relation.⁸³ But, if we do that, mathematical statements become “relativized” to the appropriate equivalence relations, which constitute additional information that needs to be carried along and managed. The alternative is to extensionalize: then the only equivalence relation one has to worry about is equality.

We do not know the extent to which Frege was familiar with examples like these. But Wilson [103, 104] calls attention to an important example of abstraction with which Frege was quite well acquainted. Frege was trained as a geometer, and studied under Ernst Schering in Göttingen. His dissertation, completed in 1893, was titled “Über eine geometrische Darstellung der imaginären Gebilde in der Ebene” (“On a geometric representation of imaginary forms in the plane”). Early nineteenth century geometers found great explanatory value and simplification in extending the usual Euclidean plane with various ideal objects, like “points at infinity” and “imaginary” points of intersection. One of the few motivating examples that Frege provided in the *Grundlagen* (§64–§68) is the fact that one can identify the “direction” of a line a in the plane with the extension of the concept “parallel to a .” As Wilson points out (though Frege does not), these “directions” are exactly what is needed to serve as points at infinity, enabling one to embed the Euclidean plane in the larger projective plane, which has a number of pleasing properties. In the projective plane, all points have equal standing, and so it stands to reason that the concept-extensions used to introduce the new entities should be given the same ontological rights as the Euclidean points and lines used in their construction. Wilson characterizes such strategies for expansion as forms of “relative logicism,” since they provide a powerful means of relating the newly-minted objects to the more familiar ones.

When it comes down to the nitty-gritty details, however, the only sustained formal development we have from Frege is his treatment of arithmetic. But even in this particular case, many of the issues we have raised come to the fore. In

⁸¹See Avigad [2], especially page 172, and Edwards [31].

⁸²See the detailed discussion in Schlimm [93, Section 3]. Other nice examples of pieces of nineteenth century mathematics that push in favor of set-theoretic abstraction are discussed in Wilson [103, 104].

⁸³Again, see Schlimm [93, Section 3].

the *Grundgesetze*, Frege defined a number of general operations and relations on tuples, sequences, functions, and relations. All of these now can be viewed as general set-theoretic constructions. What gives these constructions universal validity is that they can be applied to any domain of objects, and we now have great latitude in creating objects, as they are needed, to populate these domains. It is precisely the ability to bring a wide variety of mathematical constructions into the realm of objects, and the ability to define predicates and operations uniformly on this realm, that renders Frege’s logic so powerful—too powerful, alas. But given Frege’s goals, it should be clear why the extension operator held so much appeal.⁸⁴

To sum up, we have traced a central tension in Frege’s work to the need to balance two competing desiderata:

1. the need for flexible but rigorous ways of talking about higher-type objects, like functions, predicates, and relations, without falling prey to incoherence; and
2. the need for ways of dealing with mathematical objects uniformly, since mathematical constructions and operations have to be applied to many sorts of objects, many of which cannot be foreseen in advance.

As we have already noted, Frege is often faulted for missing the simple inconsistency that arises from the formal means he introduced to resolve this tension. Nonetheless, it is worth highlighting the extent to which these two concerns were central to the subsequent development logic and foundations. Russell’s paradox shows that Frege was perfectly right to worry that an overly naive treatment of functions, concepts, and objects would lead to problems in the most fundamental use of our language and methods of reasoning. And, going into the twentieth century, developments in all branches of mathematics called for liberal means of constructing new mathematical domains and structures, as well as uniform ways of reasoning about their essential structural properties. The most fruitful and appropriate means of satisfying these needs was by no means clear at the turn of the twentieth century. Indeed, these issues were at the heart of the tumultuous foundational debates that were looming over the horizon.

⁸⁴The same uniformity is achieved in set theory is achieved by having a large universe of sets, and incorporating set-forming operations which return new elements of that universe. Russell introduced the notion of *typical ambiguity* [40, 91] to allow “polymorphic” operations defined uniformly across types, and modern interactive theorem provers based on simple type theory follow such a strategy to obtain the necessary uniformities. For example, most such systems have operations card_σ which maps a finite set of elements of type σ to its cardinality, a natural number. The systems include mechanisms that allow one to define this family of operations uniformly, once and for all, treating σ as a parameter. One can then write $\text{card } A$, and let the system infer the relevant type parameter from the type of A . This provides one means of coping with the nonuniformities that arise from a type-theoretic compartmentalization of the mathematical universe, but the difficulties that accrue to taking simple type theory as a mathematical foundation are complex; see, for example, [6].

11 Conclusions

In 1914, Felix Hausdorff published his *Grundzüge der Mengenlehre* [55]. Dedicated to Georg Cantor, the book established a modern set-theoretic foundation, and used it to support the development of point-set topology. After discussing the notion of an ordered pair, Hausdorff gave the concept of function its modern definition:

...we consider a set P of such pairs, satisfying the condition that each element of A occurs as the first element of one and only one pair p of P . In this way, each element a of A determines a unique element b , namely, the one to which it is connected in a pair $p = (a, b)$. We denote this element associated to a , which is determined by and dependent on a , by

$$b = f(a),$$

and we say that on A (i.e. for all elements of A) a *unique function* of a is defined. We view two such functions $f(a), f'(a)$ as equal when and only when the corresponding sets of pairs P, P' are equal, that is, when, for *each* a , $f(a) = f'(a)$.⁸⁵ [55, p. 33]

Work in functional analysis in the 1910's and 1920's helped to cement the modern view. In 1859, George Boole published *A Treatise on Differential Equations* [9], in which he introduced the subject as a study of “variable quantities” subject to “known” relations between their differential coefficients. That work is noteworthy for its observation that differential operators can be viewed as algebraic expressions subject to certain laws. Integration was also viewed abstractly; in the expression $y = \int \varphi(x)dx + c$,

the symbol $\int$ denotes a certain process of integration, the study of the various forms and conditions of which is, in a peculiar sense, the object of this part of the Integral Calculus. [9, p. 2]

Modern functional analysis takes the view that operations on functions like differentiation and integration can themselves be viewed as functions, defined over spaces of other functions. In 1887, Vito Volterra published a seminal paper, “Sopra le funzioni che dipendono de altre funzioni” (“On functions that depend on other functions”) [98], which adopted such a viewpoint. In 1901, Hadamard

⁸⁵“...betrachten wir eine Menge P solcher Paare, und zwar von der Beschaffenheit, daß jedes Element a von A in einem und nur einem Paare p von P als erstes Element auftritt. Jedes Element a bestimmt auf diese Weise ein und nur ein Element b , nämlich dasjenige, mit dem es zu einem Paare $p = (a, b)$ verbunden auftritt; dieses durch a bestimmte, von a abhängige, dem a zugeordnete Element bezeichnen wir mit

$$b = f(a)$$

und sagen, daß hiermit in A (d. h. für alle Elemente von A) eine *eindeutige Funktion* von a definiert sei. Zwei solche Funktionen $f(a), f'(a)$ sehen wir dann und nur dann als gleich an, wenn die zugehörigen Paarmengen P, P' gleich sind, wenn also, für *jedes* a , $f(a) = f'(a)$ ist.”

published *Leçons sur le calcul des variations*, which helped establish the foundations of functional analysis, and, in fact, introduced the term “functional.” In 1930, Volterra published an English translation of a series of lectures he had given at the University of Madrid in 1925 [99], which began with a lengthy discussion of the notion. After presenting particular examples, he said:

We shall therefore say that a quantity z is a *functional of the function* $x(t)$ in the interval (a, b) when it depends on all the values taken by $x(t)$ when t varies in the interval (a, b) ; or, alternatively, when a law is given by which to every function $x(t)$ defined within (a, b) (the independent variable within a certain function field) there can be made to correspond one and only one quantity z , perfectly determined, and we shall write

$$z = F \left[x \begin{pmatrix} b \\ t \\ a \end{pmatrix} \right].$$

This definition of a *functional* recalls especially the ordinary general definition of a *function* given by Dirichlet. [99, p. 4]

The chapter as a whole describes an outlook that is essentially the contemporary mathematical viewpoint, yet in a way that makes clear that the viewpoint was one that the mathematical community was still getting used to.⁸⁶

Thus, by 1925, the modern view of a function was firmly in place. A function could map elements of any domain to any other; one could specify a function by specifying any determinate law; functions could serve as arguments to other functions; and functions could serve as elements of algebraic structures and geometric spaces. The famous 1905 debate among the French analysts as to whether it makes sense to consider “arbitrary functions,” not given by any rule or law, was already two decades in the past.⁸⁷

Our study of the treatment of characters in number theory has focused on only one small part of the grand historical development that resulted in this modern way of thinking. Despite its narrow focus, the case study has illuminated some important factors that contributed to the modern view. In particular, we have emphasized the difficulties inherent in developing a coherent, rigorous way of treating functions as objects, as well as concerns as to how one could do so in a way that is appropriate to mathematics and preserves all the information that is essential to a rigorous argument. We have also explored some of the benefits associated with the methodological changes. Many of these accrue to the ability of the new language and notation to highlight central features and relationships of the objects in question, while suppressing details that make it harder to discern the high-level structure of a mathematical argument. We also emphasized the importance of representing mathematical concepts and objects

⁸⁶See also the Griffith Evans’ helpful introduction to the 1959 Dover reprinting of Volterra’s lectures [99].

⁸⁷The “five letters” are translated as an appendix to Moore [80], reproduced in Ewald [39], volume 2, pages 1077–1086.

in order to capture those features that are uniform across different domains of argumentation, so that these uniformities can be packaged and used in a modular way. We noted that many of these concerns, especially concerns about coherence and uniformity, are present in Frege’s logical analysis of the function concept, despite the fact that Frege’s focus was more foundational than mathematical.

Finally, we have argued that understanding and evaluating the considerations that shape mathematical language and inferential practice is an important part of studying the metaphysics of mathematics. Whether or not one deems it appropriate to maintain a sharp distinction between a priori and empirical aspects of our scientific practice, the best way to justify our ontological posits is by understanding what they do for us, and how they help us attain our scientific goals. Moreover, we can discern among these scientific goals some that are prototypically *mathematical*. One of these is the goal of maintaining an inferential practice with clear rules and norms, one that allows its practitioners to carry out, communicate, and evaluate arguments that can become exceedingly long and complex. Another is the goal of promoting efficiency of thought, leveraging whatever features we can to extend our cognitive reach and transcend our cognitive limitations. The challenge remains that of developing appropriately philosophical means of coming to terms with such “pragmatic” and “cognitive” considerations.⁸⁸ To that end, we should resist simplistic accounts: the touchstone must always be mathematical practice itself, with all its richness and complexity.

References

- [1] William Aspray and Philip Kitcher, editors. *History and Philosophy of Modern Mathematics*. University of Minnesota Press, Minneapolis, MN, 1988.
- [2] Jeremy Avigad. Methodology and metaphysics in the development of Dedekind’s theory of ideals. In [42], pages 159–186.
- [3] Jeremy Avigad. Mathematical method and proof. *Synthese*, 153:105–159, 2006.
- [4] Jeremy Avigad. Understanding proofs. In [77], pages 317–353.
- [5] Jeremy Avigad. Understanding, formal verification, and the philosophy of mathematics. *Journal of the Indian Council of Philosophical Research*, 27:161–197, 2010.
- [6] Jeremy Avigad. Type inference in mathematics. *Bulletin of the European Association for Theoretical Computer Science*, 106:78–98, 2012.
- [7] Mark Balaguer. *Platonism and Anti-Platonism in Mathematics*. Oxford University Press, New York, 1998.

⁸⁸See Avigad [3–5] and the essays in Mancosu [77] for initial attempts in that direction.

- [8] Michael Beaney, editor. *The Frege Reader*. Blackwell Publishing, Malden, MA, 1997.
- [9] George Boole. *A Treatise on Differential Equations*. Macmillan and Co., Cambridge, 1859.
- [10] John P. Burgess. Why I am not a nominalist. *Notre Dame Journal of Formal Logic*, 24:93–105, 1983.
- [11] John P. Burgess. *Fixing Frege*. Princeton University Press, Princeton, NJ, 2005.
- [12] John P. Burgess. *Mathematics, Models, and Modality*. Cambridge University Press, Cambridge, 2008.
- [13] Georg Cantor. Beiträge zur Begründung der transfiniten Mengenlehre. *Mathematische Annalen*, 46:481–512, 1895.
- [14] Rudolf Carnap. Empiricism, semantics, and ontology. *Revue Internationale de Philosophie*, 4:20–40, 1950. Reprinted in P. Benacerraf and H. Putnam, editors, *Philosophy of Mathematics*, second edition, pages 241–257, Cambridge University Press, Cambridge, 1983.
- [15] Arthur Cayley. On the theory of groups, as depending on the symbolic equation $\theta^n = 1$. *Philosophical Magazine*, 7:40–47, 1854. Reprinted in his *Collected Mathematical Papers*, volume 2, pages 123–130, Cambridge University Press, Cambridge, 1889.
- [16] Renaud Chorlay. Questions of generality as probes into 19th century analysis. To appear in K. Chemla, R. Chorlay, D. Rabouin, editors, *Handbook on Generality in Mathematics and the Sciences*.
- [17] Alonzo Church. A formulation of the simple theory of types. *Journal of Symbolic Logic*, 5:56–68, 1940.
- [18] Charles Jean de la Vallée Poussin. Démonstration simplifiée du théorème de Dirichlet sur la progression arithmétique. *Mémoires couronnés et autres mémoires publiés par L’Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique*, 53, 1895–1896.
- [19] Charles Jean de la Vallée Poussin. *Recherches analytiques sur la théorie des nombres premiers*. Hayez, Brussels, 1897.
- [20] Richard Dedekind. Über die Einführung neuer Funktionen in der Mathematik. Delivered as a *Habitationsvorlesung* in Göttingen on June 30, 1854. Reprinted in [24], volume 3, chapter LX, pages 428–438, and translated by William Ewald as “On the introduction of new functions in mathematics” in [39], volume 2, pages 754–762.

- [21] Richard Dedekind. *Stetigkeit und irrationale Zahlen*. F. Vieweg & Sohn, Braunschweig, 1872. Reprinted in [24], volume 3, chapter L, pages 315–334. Translated by Wooster Beman as “Continuity and irrational numbers” in *Essays on the Theory of Numbers*, Open Court, Chicago, 1901; reprinted by Dover, New York, 1963. The Beman translation is reprinted, with corrections by William Ewald, in [39], volume 2, pages 765–779.
- [22] Richard Dedekind. *Was sind und was sollen die Zahlen?* F. Vieweg & Sohn, Braunschweig, 1888. A later edition reprinted in [24], Volume 3, Chapter LI, pages 335–391. The second edition, with a new preface, was published in 1893, and is translated by Wooster Beman as “The nature and meaning of numbers” in *Essays on the Theory of Numbers*, Open Court, Chicago, 1901; reprinted by Dover, New York, 1963. The Beman translation is reprinted, with corrections by William Ewald, in [39], volume 2, pages 787–833.
- [23] Richard Dedekind. Über die Theorie der ganzen algebraischen Zahlen. Supplement XI to the fourth edition of [29], 1894, pages 434–657. Reprinted in [24], volume 3, chapter 46, pages 1–222.
- [24] Richard Dedekind. *Gesammelte mathematische Werke*, volumes 1–3, edited by Robert Fricke, Emmy Noether and Øystein Ore, F. Vieweg & Sohn, Braunschweig, 1932. Reprinted by Chelsea Publishing Co., New York, 1968.
- [25] Johann Peter Gustav Lejeune Dirichlet. Sur la convergence des séries trigonométriques qui servent à représenter une fonction arbitraire entre des limites données. *Journal für die reine und angewandte Mathematik*, 4:157–169, 1829. Reprinted in [30], pages 117–132.
- [26] Johann Peter Gustave Lejeune Dirichlet. Ueber eine neue anwendung bestimmter integrale auf die summation endlicher oder unendlicher reihen. *Abhandlungen der königlich Preussischen Akademie der Wissenschaften*, pages 391–407, 1835. Reprinted in [30], pages 237–256.
- [27] Johann Peter Gustav Lejeune Dirichlet. Beweis eines Satzes über die arithmetische Progression. *Bericht über die Verhandlungen der königlich Preussischen Akademie der Wissenschaften Berlin*, 1837. Reprinted in [30], pages 307–312.
- [28] Johann Peter Gustav Lejeune Dirichlet Dirichlet. Beweis des Satzes, dass jede unbegrenzte arithmetische Progression, deren erstes Glied und Differenz ganze Zahlen ohne gemeinschaftlichen Factor sind, unendlich viele Primzahlen enthält. *Abhandlungen der königlich Preussischen Akademie der Wissenschaften*, pages 45–81, 1837. Reprinted in [30], pg 313–342. Translated by Ralf Stefan as “There are infinitely many prime numbers in all arithmetic progressions with first term and difference coprime,” arXiv:0808.1408.

- [29] Johann Peter Gustave Lejeune Dirichlet. *Vorlesungen über Zahlentheorie*. Vieweg, Braunschweig, 1863. Edited by Richard Dedekind. Subsequent editions in 1871, 1879, 1894, with “supplements” by Richard Dedekind. Translated by John Stillwell, with introductory notes, as *Lectures on Number Theory*, American Mathematical Society, Providence, RI, 1999.
- [30] Johann Peter Gustav Lejeune Dirichlet. *Werke*, edited by Leopold Kronecker. Georg Reimer, Berlin, 1889.
- [31] Harold M. Edwards. The genesis of ideal theory. *Archive for History of Exact Sciences*, 23:321–378, 1980.
- [32] Harold M. Edwards. Kronecker’s views on the foundations of mathematics. In D. E. Rowe and J. McCleary, editors, *The History of Modern Mathematics*, pages 67–77, Academic Press, San Diego, 1989.
- [33] Harold M. Edwards. Kronecker’s fundamental theorem of general arithmetic. In Jeremy Gray and Karen Parshall, editors, *Episodes in the History of Modern Algebra (1800–1950)*, pages 107–116, American Mathematical Society, Providence, RI, 2007.
- [34] Harold M. Edwards. Kronecker’s algorithmic mathematics. *Mathematical Intelligencer*, 31:11–14, 2009.
- [35] Gotthold Eistenstein. Eine neue Gattung zahlentheoretischer Funktionen, weche von zwei Elementen abhängen und durch gewisse lineare Funktional-Gleichungen definirt werden. *Bericht über die zur Bekanntmachung geeigneten Verhandlungen der königlich Preussischen Akademie der Wissenschaften zu Berlin*, pages 36–42, 1850. Reprinted in Eisenstein’s *Mathematische Werke*, volume 2, pages 705–711, Chelsea Publishing Company, New York, 1989.
- [36] Leonhard Euler. *Introductio in analysin infinitorum, tomus primus*. Lausannae, 1748. Publications E101 and E102 in the Euler Archive.
- [37] Leonhard Euler. Speculationes circa quasdam insignes proprietates numerorum. *Acta Academiae Scientiarum Imperialis Petropolitinae*, 4, 1784. Publication E564 in the Euler Archive. Translated by Jordan Bell as “Speculations on some characteristic properties of numbers,” 2007, arXiv:0705.3929.
- [38] Graham Everest and Thomas Ward. *An Introduction to Number Theory*. Springer, London, 2005.
- [39] William Ewald, editor. *From Kant to Hilbert: A Source Book in the Foundations of Mathematics*, volumes 1 and 2. Oxford University Press, Oxford, 1996.
- [40] Solomon Feferman. Typical ambiguity: trying to have your cake and eat it too. In Godehard Link, editor, *One hundred years of Russell’s paradox*, pages 135–151, de Gruyter, Berlin, 2004.

- [41] José Ferreirós. *Labyrinth of Thought: A History of Set Theory and its Role in Modern Mathematics*. Birkhäuser, Basel, 1999.
- [42] José Ferreirós and Jeremy Gray, editors. *The Architecture of Modern Mathematics*. Oxford University Press, Oxford, 2006.
- [43] Gottlob Frege. *Begriffsschrift, eine der arithmetischen nachgebildete Formelsprache des reinen Denkens*. Louis Nebert, Halle, 1879. Translated by Michael Beaney in [8], pages 84–129.
- [44] Gottlob Frege. *Die Grundlagen der Arithmetik*. W. Koebner, Breslau, 1884. Translated by J. L. Austin as *The Foundations of Arithmetic*, second revised edition, Northwestern University Press, Evanston, Illinois, 1980. Excerpts translated by Michael Beaney in [8], pages 84–129.
- [45] Gottlob Frege. *Function und Begriff*. Hermann Pohle, Jena, 1891. Reprinted in [49], and translated by Peter Geach as “Function and concept” in [8]. Page number references are to the original publication.
- [46] Gottlob Frege. Über Begriff und Gegenstand. *Vierteljahresschrift für wissenschaftliche Philosophie*, 16:192–505, 1892. Reprinted in [49] and translated by Peter Geach as “Concept and object” in [8]. Page number references are to the original publication.
- [47] Gottlob Frege. *Grundgesetze der arithmetik*. H. Pohle, Jena, volume 1, 1893, volume 2, 1903. Excerpts translated by Michael Beaney in [8], pages 194–223 and 258–289.
- [48] Gottlob Frege. Was ist eine Funktion? In Stefan Meyer, editor, *Festschrift Ludwig Boltzmann gewidmet zum sechzigsten Geburtstage*. J. A. Barth, Leipzig, 1904. Reprinted in [49] and translated as “What is a function?” in Peter Geach and Max Black, *Translations from the Philosophical Writings of Gottlob Frege*, Oxford University Press, Oxford, 1980. Page number references are to the original publication.
- [49] Gottlob Frege. *Funktion – Begriff – Bedeutung*, edited by Mark Textor. Vandenhoeck and Ruprecht, Göttingen, 2002.
- [50] Carl Friedrich Gauss. *Disquisitiones Arithmeticae*. G. Fleischer, Leipzig, 1801. Reprinted in Gauss’ *Werke*, Königlichen Gesellschaft der Wissenschaften, Göttingen, 1863. Translated with a preface by Arthur A. Clarke, Yale University Press, New Haven, 1966, and republished by Springer, New York, 1986.
- [51] M. J. C. Gordon and T. F. Melham, editors. *Introduction to HOL: A Theorem Proving Environment for Higher-Order Logic*. Cambridge University Press, Cambridge, 1993.
- [52] Umberto Bottazzini and Jeremy Gray. *Hidden Harmony—Geometric Fantasies: The Rise of Complex Function Theory*. Springer, Berlin, 2013.

- [53] Jacques Hadamard. Sur la distribution des zéros de la fonction $\zeta(s)$ et ses conséquences arithmétiques. *Bulletin de la Société Mathématique de France*, 24:199–220, 1896.
- [54] John Harrison. HOL light: a tutorial introduction. In Mandayam Srivas and Albert Camilleri, editors, *Proceedings of the First International Conference on Formal Methods in Computer-Aided Design*, pages 265–269, Springer, Berlin, 1996.
- [55] Felix Hausdorff. *Grundzüge der Mengenlehre*. Veit and Company, Leipzig, 1914. Reprinted by Chelsea Publishing Company, New York, 1949.
- [56] Thomas Hawkins. The origins of the theory of group characters. *Archive for History of Exact Sciences*, 7:142–170, 1971.
- [57] Richard Heck. The Julius Ceasar objection. In Richard Heck, editor, *Language, Thought, and Logic: Essays in Honour of Michael Dummett*, pages 273–308, Oxford University Press, Oxford, 1997. Reprinted in Richard Heck, editor, *Frege’s Theorem*, pages 127–155, Oxford University Press, Oxford, 2011.
- [58] Richard Heck and Robert May. The composition of thoughts. *Noûs*, 45:126–166, 2011.
- [59] Philip Kitcher. *The Nature of Mathematical Knowledge*. Oxford University Press, New York, 1983.
- [60] Philip Kitcher. Mathematical naturalism. In [1], pages 293–325.
- [61] Felix Klein. A comparative review of recent researches in geometry. *Bulletin of the American Mathematical Society*, 2:215–249, 1893.
- [62] Israel Kleiner. Evolution of the function concept: a brief survey. *The College Mathematical Journal*, 20:282–300, 1989.
- [63] Leopold Kronecker. Auseinandersetzung einiger eigenschaften der klassenzahl idealer complexer zahlen. *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin*, pages 881–882, 1870. Reproduced in [67], volume I, pages 271–282.
- [64] Leopold Kronecker. *Grundzüge einer arithmetischen Theorie der algebraischen Grössen*. Riemeier, Berlin, 1882. Also published in *Journal für reine und angewandte Mathematik*, volume 92, 1882, pages 1–122, and [67], volume II, pages 237–387.
- [65] Leopold Kronecker. Ein Fundamentalsatz der allgemeinen Arithmetik. *Journal für die reine und angewandte Mathematik*, 100:490–510, 1887. Reprinted in [67], volume IIIa, pages 209–240.
- [66] Leopold Kronecker. *Vorlesungen über Zahlentheorie*, edited by Kurt Hensel, B. G. Teubner, Leipzig, 1901. Republished by Springer, Berlin, 1978.

- [67] Leopold Kronecker. *Werke*, edited by Kurt Hensel, volumes 1–5, B. G. Teubner, Leipzig, 1895–1930. Reprinted by Chelsea Publishing Co., New York, 1968.
- [68] Edmund Landau. *Handbuch der Lehre von der Verteilung der Primzahlen*, volume 1. B. G. Teubner, Leipzig, 1909.
- [69] Edmund Landau. *Vorlesungen über Zahlentheorie*. S. Hirzel, Leipzig, 1927. Translated by Jacob E. Goodman as *Elementary Number Theory*, Chelsea Publishing Company, New York, 1958.
- [70] Adrien-Marie Legendre. Recherches d’analyse indéterminée. *Histoire de l’Académie Royale des Sciences de Paris*, pages 465–559, 1788 (1785).
- [71] Mary Leng. *Mathematics and Reality*. Oxford University Press, Oxford, 2010.
- [72] Nikolai Luzin. Functions. *The American Mathematical Monthly*, 105:59–67 (Part I) and 105:263–270 (Part II), 1998.
- [73] Ernst Mach. *The Science of Mechanics: A Critical and Historical Account of its Development*, translated by Thomas J. McCormack. The Open Court Publishing Co., La Salle, Illinois, 1960.
- [74] George W. Mackey. Harmonic analysis as the exploitation of symmetry: a historical survey. *Bulletin of the American Mathematical Society*, 3:543–698, 1980.
- [75] Penelope Maddy. *Naturalism in Mathematics*. Oxford University Press, New York, 1997.
- [76] Penelope Maddy. *Second Philosophy: A Naturalistic Method*. Oxford University Press, Oxford, 2007.
- [77] Paolo Mancosu, editor. *The Philosophy of Mathematical Practice*. Oxford University Press, Oxford, 2008.
- [78] Kenneth Manders. The Euclidean diagram. In [77], pages 80–133.
- [79] A.F. Monna. The concept of function in the 19th and 20th centuries, in particular with regard to the discussions between Baire, Borel and Lebesgue. *Archive for History of Exact Sciences*, 9:57–84, 1972.
- [80] Gregory H. Moore. *Zermelo’s Axiom of Choice: Its Origins, Development, and Influence*. Springer, New York, 1982.
- [81] Rebecca Morris. *Character and object*. Master’s thesis, Carnegie Mellon University, 2011.
- [82] Tobias Nipkow, Lawrence C. Paulson, and Markus Wenzel. *Isabelle/HOL. A Proof Assistant for Higher-Order Logic*. Springer, Berlin, 2002.

- [83] Marco Panza. From Lagrange to Frege: functions and expressions. In Carlos Alvarez and Andrew Arana, editors, *Analytic Philosophy and the Foundations of Mathematics*, Palgrave Macmillan, to appear.
- [84] David Pengelley. Arthur Cayley and the first paper on group theory. In Amy Shell-Gellasch and Dick Jardine, editors, *From Calculus to Computers: Using the Last 200 Years of Mathematics History in the Classroom*, pages 3–8, Mathematics Association of America, Washington, D.C., 2005.
- [85] W. V. O. Quine. Main Trends in Recent Philosophy: Two Dogmas of Empiricism. *The Philosophical Review*, 60:20–43, 1951.
- [86] W. V. O. Quine. On Frege’s way out. *Mind*, 64:145–159, 1955.
- [87] W. V. O. Quine. *Ontological Relativity, and other Essays*. Columbia University Press, New York, 1969.
- [88] W. V. O. Quine. *Word and Object*. MIT Press, Cambridge, MA, 1960.
- [89] W. V. O. Quine. On what there is. *The Review of Metaphysics*, 2:21–38, 1948. Reprinted in W. V. O. Quine, *From a Logical Point of View*, Harvard University Press, Cambridge, 1980.
- [90] Erich Reck. Frege-Russell numbers: analysis or explication? In Michael Beaney, editor, *The Analytic Turn: Analysis in Early Analytic Philosophy and Phenomenology*, pages 33–50, Routledge, London, 2007.
- [91] Bertrand Russell. Mathematical logic as based on the theory of types. *American Journal of Mathematics*, 30:222–262, 1908.
- [92] Bertrand Russell and Alfred North Whitehead. *Principia Mathematica*. Cambridge University Press, Cambridge, first volume, 1910; second volume, 1912; third volume, 1913.
- [93] Dirk Schlimm. On abstraction and the importance of asking the right research questions: could Jordan have proved the Jordan-Hölder theorem? *Erkenntnis*, 68:409–420, 2008.
- [94] Howard Stein. Logos, logic, and logistiké. In [1], pages 238–259.
- [95] Jamie Tappenden. The Riemannian background to Frege’s philosophy. In [42], pages 97–132.
- [96] A. S. Troelstra and Dirk van Dalen. *Constructivism in Mathematics: An Introduction*, volumes 1 and 2. North-Holland, Amsterdam, 1988.
- [97] Alasdair Urquhart. Mathematics and physics: strategies of assimilation. In [77], pages 417–440.
- [98] Vito Volterra. Sopra le funzioni che dipendono de altre funzioni. *Rend. R. Accademia dei Lincei*, 2:97–105, 141–146 and 153–158, 1887.

- [99] Vito Volterra. *Theory of Functionals and of Integral and Integro-differential Equations*. Blackie and Son, Ltd., London and Glasgow, 1930. Translated by M. Long and edited by Luigi Fantappiè. Reprinted with a preface by G. C. Evans, a biography of Vito Volterra and a bibliography of his published works by E. Whittaker, Dover, New York, 1959.
- [100] Heinrich Weber. Beweis des Satzes, dass jede eigentlich primitive quadratische Form unendlich viele Primzahlen darzustellen fähig ist. *Mathematische Annalen*, 20:301–329, 1882.
- [101] Mark Wilson. Can we trust logical form? *The Journal of Philosophy*, 91:519–544, 1994.
- [102] Mark Wilson. *Wandering Significance: An Essay on Conceptual Behavior*. Oxford University Press, Oxford, 2006.
- [103] Mark Wilson. Frege’s mathematical setting. In Michael Potter and Tom Ricketts, editors, *The Cambridge Companion to Frege*, Cambridge University Press, Cambridge, 2010.
- [104] Mark Wilson. Enlarging one’s stall *or* How did all of these *sets* get in here? Preprint.
- [105] Ludwig Wittgenstein. *Wittgenstein’s Lectures on the Foundations of Mathematics, Cambridge, 1939*. University of Chicago Press, Chicago, 1989.
- [106] Hans Wussing. *The Genesis of the Abstract Group Concept: A Contribution to the History of the Origin of Abstract Group Theory*, translated from the German by Abe Shenitzer and Hardy Grant. MIT Press, Cambridge, MA, 1984.
- [107] A. P. Youschkevitch. The concept of function up to the middle of the 19th century. *Archive for History of Exact Sciences*, 16:37–85, 1976.